

THE MACROSCOPICAL SOUND OF TORI

by Constantin Vernicos

Abstract

Take a torus with a Riemannian metric. Lift the metric on its universal cover. You get a distance which in turn yields balls. On these balls you can look at the Laplacian. Focus on the spectrum for the Dirichlet or Neumann problem. We describe the asymptotic behaviour of the eigenvalues as the radius of the balls goes to infinity, and characterise the flat tori using the tools of homogenisation our conclusion being that "Macroscopically, one can hear the shape of a flat torus". We also show how in the two dimensional case we can recover earlier results by D. Burago and S. Ivanov on the asymptotic volume.

1 Introduction and claims

Let $(\mathbb{T}^n, g)$ be a riemannian Torus, lift its metric on its universal cover and use it to define first a distance, then the metric's balls. The first thing one can observe is the volume of these balls as a function of their radius, indeed as the distance obtained arises from a compact quotient it is equivalent to an euclidean distance hence the volume of these balls is equivalent to the euclidean volume of an euclidean ball i.e. proportional to the radius of the ball to the power of n (the dimension of our Torus).

We are thus naturally led to wonder what happens if one looks at the following riemannian function on the balls ($B_g(\rho)$ is the ball of radius ρ)

$$\frac{\text{Vol}_g(B_g(\rho))}{\rho^n} \quad \text{as } \rho \rightarrow +\infty .$$

If it is not very surprising that it converges to some constant for this limit can be seen as a mean value due to the periodicity of the metric (see for example Pansu [Pan82] and a slightly different and more analytical proof in this paper section 2.3), it is quite remarkable that this constant, called asymptotic volume, is bounded from below by the constant arising from the flat tori and furthermore that the case of equality characterises the flat tori as D. Burago and S. Ivanov showed in [BI95].

The study of the balls of large radii on the universal cover of tori (and more generally of a nilmanifold) is what we call here the macroscopical geometry. Indeed in our case the universal cover is a real vector space, where some lattices acts by

Key-words : Spectrum of the laplacian, Tori, Homogenisation, Stable norm, Asymptotic volume.

Classification : 53C24, 58C40, 74Q99

translation (in the more general case of nilmanifolds one should consider a left action). Should one focus on the point of this lattice endowed with the distance arising from the torus, one gets an invariant metric on the lattice. Now if one looks at this lattice from a galaxy far, far away, one won't be able to distinguish the lattice from the whole universal cover. Thus it is understandable that for this observer the distance obtained on the universal cover seems invariant by all translations (for general nilmanifolds one gets a left invariant distance).

In the case of tori this "seen from a far away galaxy" distance is a norm, called the stable norm and was first defined by Federer in Homology. It is some kind of mean value of the metric. This asymptotic behaviour was generalized and proved by P. Pansu for all nilmanifold [Pan82] and precised by D. Burago [Bur92] for Tori. Since then the stable norm appeared in many other works : for surfaces and the links with Aubry-Mather theory in D. Massart's works, one can also find it in the weak KAM theory of A. Fathi. It is also worth mentioning the crucial role it plays in the proof by D. Burago and S. Ivanov [BI94] of the Hopf conjecture concerning tori without conjugate points. Here in part 2 we show, for the case of tori, how one recovers the stable norm using Homogenisation's tools.

There is another interesting geometric invariant attached to the balls and linked with the volume, the spectrum of the laplacian. Indeed if one knows the spectrum one knows the volume thanks to Weyl's asymptotic formula. Here again one easily sees, comparing with the euclidean case, that the eigenvalues converge to zero with a $1/\rho^2$ speed (ρ being the radius). If one can expect a convergence when rescaled, it is quite surprising that as a limit we obtain the spectrum of an euclidean and not a finslar metric, indeed the behaviour is described by the following theorem which is one of the aims of this paper :

Theorem 1. *Let $(\mathbb{T}^n, g)$ be a Riemannian Torus, $B_g(\rho)$ the induced metric ball on its universal cover and $\lambda_i(B_g(\rho))$ the i^{th} eigenvalue of the Laplacian for the Dirichlet (resp. Neumann) problem.*

There exists an elliptic operator Δ_∞ , which is the Laplacian of some Euclidean metric on $\mathbb{R}^n$, such that if λ_i^∞ is its i^{th} eigenvalue for the Dirichlet (resp. Neumann) problem on the stable's norm unit ball then

$$\lim_{\rho \rightarrow +\infty} \rho^2 \lambda_i(B_g(\rho)) = \lambda_i^\infty$$

The section 4 is devoted to the proof of this theorem and section 3 introduces the analytical background : Homogenisation and various convergence fairly known by the specialist of homogenisation but adapted here to our purpose, as it becomes after some re-spelling of the problem made in section 2.4. From this theorem we can deduce the following one which is some kind of "à la" Burago-Ivanov macroscopical rigidity and which inspired the abstract :

Theorem 2. *Let $(\mathbb{T}^n, g)$ be a Riemannian Torus, $B_g(\rho)$ the induced metric ball on its universal cover and $\lambda_1(B_g(\rho))$ the first eigenvalue of the Laplacian for the Dirichlet problem. Then*

$$(a) \lim_{\rho \rightarrow +\infty} \rho^2 \lambda_1(B_g(\rho)) = \lambda_1^\infty \leq \lambda_{e,n}$$

(b) equality holds if, and only if, the torus is flat

where $\lambda_{e,n}$ is the first eigenvalue of the Euclidean Laplacian on the Euclidean unit ball.

The proof which is done in section 6, involves some kind of transplantation for the inequality mixed with Γ -convergence for the equality. For a better understanding of what happens we briefly give some informations related to the Γ -convergence and adapt it to our purpose in section 5, following the general ideas of K. Kuwae and T. Shioya in [KS] (who in turn generalized U. Mosco's paper [Mos94]), this section being completed by the proof of section 8.

As the macroscopical spectrum involved rises from an euclidean metric, we can use the Faber-Krahn inequality to obtain a new inequality regarding the asymptotic volume, this is done in section 7.1 :

Proposition 3. *Let $(\mathbb{T}^n, g)$ be a Riemannian Torus, $B_g(\rho)$ the geodesic balls of radius ρ centred on a fixed point and $\text{Vol}_g(B_g(\rho))$ their Riemannian volume induced on the universal cover, writing*

$$\text{Asvol}(g) = \lim_{\rho \rightarrow \infty} \frac{\text{Vol}_g(B_g(\rho))}{\rho^n}$$

then

1. $\text{Asvol}(g) \geq \frac{\text{Vol}_g(\mathbb{T}^n)}{\text{Vol}_{\text{Al}}(\mathbb{T}^n)} \omega_n$
2. In case of equality, the torus is flat.

Where ω_n is the unit Euclidean ball's Euclidean volume, and $\text{Vol}_{\text{Al}}(\mathbb{T}^n)$ is the volume of the albanese Torus.

A surprising facts arises because this new inequality involves a constant which happened to be at the heart of the isosystolic inequality of two dimensional tori (see J. Lafontaine [Laf74]), hence we obtain an alternate proof of the asymptotic volume's lower boundedness in dimension two,

Corollary. *Let $(\mathbb{T}^2, g)$ be a 2-dimensional torus then :*

1. $\text{Asvol}(g) \geq \pi$
2. In case of equality, the torus is flat.

It is worth mentioning that the case of equality in the previous claims relies on theorem 32, which states that the stable norm coincides with the Albanese metric if and only if the torus is flat, and whose proof do not depend on the work of D. Burago and S. Ivanov [BI95], thus we actually get an alternate proof of their theorem in the 2-dimensional case.

We also give a kind of generalised Faber-Krahn inequality for normed finite dimensional vector spaces, which implies that we cannot distinguish the euclidean's ones among them using the first generalised eigenvalue of the Dirichlet laplacian (see lemma 35 and its corollary):

Theorem 4. (Faber-Krahn inequality for norms). *Let D be a domain of $\mathbb{R}^n$, with the norm $\|\cdot\|$ and a measure μ invariant by translation. Let D^* be the norm's ball with same measure as D , then*

$$\lambda_1(D^*, \|\cdot\|) \leq \lambda_1(D, \|\cdot\|)$$

We finally explain in section 7.2 how is our work related to works focused on the long time asymptotics of the heat kernel (see [KS00], [DZ00], [ZKON79]) and finally in section 7.3 we state how theorem 1 transposes to all graded nilmanifolds (subject which should be widely extended in a forthcoming article).

I would like to thanks my advisor Gérard Besson for his good advice and Y. Colin de Verdière who lead me to the Γ -convergence which happened to be the adapted tool, as I hope to convince the reader, for the macroscopical geometry.

2 Stable Norm and Homogenisation

In this section we show how the stable norm, the Gromov-Hausdorff convergence and the Γ -convergence of the homogenisation theory are linked and finish by re-spelling our goal. In what follows, $B_g(\rho)$ will be the metric ball of radius ρ on the universal cover of a torus with the lifted metric. We first begin by two definitions.

2.1 Convergences

We recall the definition of Γ -convergence in a metric space :

Definition 5. *Let (X, d) be a metrics space. We say that a sequence of function (F_j) from X to $\overline{\mathbb{R}}$, Γ -converges to a function $F : X \rightarrow \overline{\mathbb{R}}$ if and only if for all $x \in X$ we have*

1. *for all converging sequence (x_j) to x*

$$F(x) \leq \liminf_{j \rightarrow \infty} F_j(x_j);$$

2. *there exists a sequence (x_j) converging to x such that*

$$F(x) = \lim_{j \rightarrow \infty} F_j(x_j).$$

We now introduce the Gromov-Hausdorff measured convergence in the space $\mathcal{M}$ of compact metric and measured spaces (X, d, m) modulo isometries. First if X and Y are in $\mathcal{M}$ then an application $\phi : X \rightarrow Y$ is called an ϵ -Hausdorff approximation if and only if we have

- (1.) the ϵ -neighbourhood of $\phi(X)$ in Y is Y ;
- (2.) for all $x, y \in X$ we have

$$\left| d(x, y) - d(\phi(x), \phi(y)) \right| \leq \epsilon.$$

We write $C^0(X)$ for the space of continues functions from X to $\mathbb{R}$ and $\mathcal{A}$ will be a partially directed space.

Definition 6. We say that a net $(X_\alpha, d_\alpha, m_\alpha)_{\alpha \in \mathcal{A}}$ of spaces in $\mathcal{M}$ converges to (X, d, m) for the Gromov-Hausdorff measured topology if, and only if there exists a net of positive real numbers $(\varepsilon_\alpha)_{\alpha \in \mathcal{A}}$ decreasing to 0 and m_α measurable ε_α -Hausdorff approximations $f_\alpha : X_\alpha \rightarrow X$ such that $(f_\alpha)_*(m_\alpha)$ converges vaguely to m i.e.

$$\int_{X_\alpha} u \circ f_\alpha \, dm_\alpha \rightarrow \int_X u \, dm \quad \forall u \in C^0(X).$$

2.2 The stable norm

Let $(\mathbb{T}^n, g)$ a Riemannian torus. We will call *rescaled metrics* the metrics $g_\rho = (1/\rho^2)(\delta_\rho)^*g$ and their lifts on the universal cover. We will also write δ_ρ for the homothetic of scale ρ .

In the 80's P. Pansu showed that the distance induced on $\mathbb{R}^n$ as a universal cover of a torus, behaved asymptotically like the distance induced by a norm. In the 90's D. Burago showed a similar result for periodic metrics on $\mathbb{R}^n$. It is that norm which is called the *stable norm*. To be more precise let us write $f_1(x) = d_g(0, x)$ the distance from the origin to x and $f_\rho(x) = d_g(0, \delta_\rho(x))/\rho$, then P. Pansu's result says that there exist a norm $\|\cdot\|_\infty$ such that for all $x \in \mathbb{R}^n$

$$\lim_{\rho \rightarrow \infty} f_\rho(x) = \|x\|_\infty$$

and Burago's says that there exists a constant C such that for all $x \in \mathbb{R}^n$

$$|f_\rho(x) - \|x\|_\infty| \leq \frac{C}{\rho}$$

in other words, Pansu's results is a simple convergence and Burago's is a uniform convergence result.

There is another proof of the simple convergence of the sequence (f_ρ) as ρ goes to infinity, using homogenisation's tools.

Theorem 7. Let $\tilde{g}$ the induced metric on $\mathbb{R}^n$ as a universal cover of a Riemannian Torus $(\mathbb{T}^n, g)$, then there exists a norm $\|\cdot\|_\infty$ such that

1. for every bounded open $I \subset \mathbb{R}$ the sequence of functional on $W^{1,2}(I; \mathbb{R}^n)$

$$E_\rho(u) = \int_I \tilde{g}_{(\delta_\rho u(t))}(u'(t), u'(t)) dt$$

Γ -converge for the L^2 norm toward the functional

$$E_\infty(u) = \int_I \|u'(t)\|_\infty^2 dt;$$

2. the norm satisfies

$$\|\xi\|_\infty = \lim_{t \rightarrow +\infty} \inf \left\{ \frac{1}{t} \int_0^t \tilde{g}_{(u+\xi\tau)}(u' + \xi, u' + \xi) \, d\tau : u \in W_0^{1,2}([0, t[; \mathbb{R}^n) \right\}. \quad (1)$$

Furthermore if $f_\rho(x) = d_g(0, \rho x)/\rho$ then for all $x \in \mathbb{R}^n$

$$\lim_{\rho \rightarrow +\infty} f_\rho(x) = \|x\|_\infty$$

Proof. We use the Proposition 16.1 page 142 of A. Braides and A. Desfranceschi . It gives us the Γ -convergence of the sequence of functional (E_ρ) toward a functional E_∞ such that

$$E_\infty(u) = \int_I \varphi(u'(t)) dt$$

with φ convex and satisfying the asymptotic formula (1). It remains to show that φ is the square of a norm.

Homogeneity : Using the asymptotic formula (1) we easily get $\varphi(0) = 0$ and by a change of variables $\varphi(\lambda x) = \lambda^2 \varphi(x)$.

Separation : Let us point that

1. The minimum of the energy of a path between 0 and $t\xi$ in an Euclidean space is attained for the straight line. Thus if we put into the asymptotic formula (1) an Euclidean metric, we get the same metric.
2. Let g and h be two metrics such that for all s and ξ

$$g_s(\xi, \xi) \leq h_s(\xi, \xi)$$

then for all $u \in W_0^{1,2}([0, t[; \mathbb{R}^n)$ we get

$$\frac{1}{t} \int_0^t g_{(u+\xi\tau)}(u' + \xi, u' + \xi) d\tau \leq \frac{1}{t} \int_0^t h_{(u+\xi\tau)}(u' + \xi, u' + \xi) d\tau$$

thus taking the infimum for u and taking the limit as t goes to infinity we get

$$\begin{aligned} & \lim_{t \rightarrow +\infty} \inf \left\{ \frac{1}{t} \int_0^t g_{(u+\xi\tau)}(u' + \xi, u' + \xi) d\tau : u \in W_0^{1,2}([0, t[; \mathbb{R}^n) \right\} \leq \\ & \lim_{t \rightarrow +\infty} \inf \left\{ \frac{1}{t} \int_0^t h_{(u+\xi\tau)}(u' + \xi, u' + \xi) d\tau : u \in W_0^{1,2}([0, t[; \mathbb{R}^n) \right\}. \end{aligned} \quad (2)$$

Now let us also remark that g being periodic, there exists two strictly positive constants α and β such that

$$\alpha|\xi|^2 \leq g_s(\xi, \xi) \leq \beta|\xi|^2$$

now applying the three remarks we get

$$\alpha|\xi|^2 \leq \varphi(\xi) \leq \beta|\xi|^2$$

thus $\varphi(\xi) = 0$ if and only if $\xi = 0$.

Triangle inequality : First note that

$$\{\xi \in \mathbb{R}^n \mid \varphi(\xi) \leq 1\} = \{\xi \in \mathbb{R}^n \mid \sqrt{\varphi(\xi)} \leq 1\} = S_n$$

It follows that if $\sqrt{\varphi}(\xi) = 1 = \varphi(\xi)$ and $\sqrt{\varphi}(\eta) = 1 = \varphi(\eta)$ then for all $0 \leq \lambda \leq 1$ by the convexity of φ

$$\varphi(\lambda\xi + (1-\lambda)\eta) \leq \lambda\varphi(\xi) + (1-\lambda)\varphi(\eta) = 1$$

so

$$\sqrt{\varphi}(\lambda\xi + (1-\lambda)\eta) \leq 1.$$

Thus for all non null x, y

$$\sqrt{\varphi}\left(\lambda\frac{x}{\sqrt{\varphi}(x)} + (1-\lambda)\frac{y}{\sqrt{\varphi}(y)}\right) \leq 1$$

now taking $\lambda = \sqrt{\varphi}(x)/(\sqrt{\varphi}(x) + \sqrt{\varphi}(y))$ and using $\sqrt{\varphi}$ homogeneity we finally get the triangle inequality and we are able to conclude that $\|\cdot\|_\infty = \sqrt{\varphi}(\cdot)$ is a norm.

The final assertion comes from the fact that $\|\xi\|_\infty^2$ is the limit of the energies' infimum along the paths between 0 and ξ for the rescaled metrics $(1/t^2)(\delta_t^*)g$, which are attained along the geodesics. $\square$

This theorem easily induces the following assertion.

Corollary 8. *For all x and $y \in \mathbb{R}^n$ we have*

$$\lim_{\rho \rightarrow +\infty} \frac{d_g(\rho x, \rho y)}{\rho} = \|x - y\|_\infty$$

From now on we will write $d_\rho(x, y) = d_g(\rho x, \rho y)/\rho$, and we are now going to see what can be deduced for the balls $B_g(\rho)$ in terms of Gromov-Hausdorff convergence.

2.3 Gromov-Hausdorff convergence of metric balls

We will write μ_g (resp. μ_ρ) the measure induced by $\tilde{g}$ (resp. g_ρ). μ_∞ will be the measure of Lebesgue such that for a fundamental domain D_f we have $\mu_\infty(D_f) = \mu_g(D_f)$. Finally let

$$B_\rho(R) = \{x \in \mathbb{R}^n \mid d_\rho(0, x) \leq R\} = \frac{1}{\rho} B_g(R \cdot \rho),$$

and

$$B_\infty(R) = \{x \in \mathbb{R}^n \mid \|x\|_\infty \leq R\}.$$

Theorem 9. *The net of measured metric spaces $(B_\rho(1), d_\rho, \mu_\rho)$ converges in the Gromov-Hausdorff measured topology to $(B_\infty(1), \|\cdot\|_\infty, \mu_\infty)$ as ρ goes to infinity.*

Proof. Let us choose an $\epsilon > 0$. We first show that the identity is an ϵ -approximation if ρ is large enough. It suffice to show that there is a finite family of points $(x_1, \dots, x_N)$ such that its ϵ -neighbourhood in $(B_\infty(1), d_\infty)$ and, for ρ large enough, in $(B_\rho(1), d_\rho)$ is respectively $B_\infty(1)$ and $B_\rho(1)$ and such that for all $i, j = 1, \dots, N$ we have

$$\left| \|x_i - x_j\|_\infty - d_\rho(x_i, x_j) \right| \leq \epsilon$$

Let $r > 0$ and let $(\gamma_1, \dots, \gamma_N)$ be all the images of 0 by the action of $\mathbb{Z}^n$, such that for $i = 1, \dots, N$, $\gamma_i \in B_\infty(r)$. Then we take for $i = 1, \dots, N$, $x_i = \gamma_i/r$. Let us remark that for ρ large enough these points will all be in $B_\rho(1)$.

Now let us point out that, because of the invariance by the $\mathbb{Z}^n$ action, there are two constants α and β such that for all x and $y \in \mathbb{R}^n$ we have

$$\alpha \|x - y\|_\infty \leq d_g(x, y) \leq \beta \|x - y\|_\infty;$$

thus for every $x \in B_\infty(1)$ take the closest point x_i (thus γ_i is the closest point of $\mathbb{Z}^n \cdot 0$ from rx) then there is a constant C (the diameter of the fundamental domain) such that

$$\|x - x_i\|_\infty \leq \frac{1}{\alpha r} d_g(rx, \gamma_i) \leq \frac{1}{\alpha r} C$$

we also get

$$d_\rho(x, x_i) \leq \frac{\beta}{\alpha r} C$$

thus, for r large enough $(x_1, \dots, x_N)$ is an ϵ -neighbourhood of $(B_\infty(1), \|\cdot\|_\infty)$. Furthermore if ρ is large enough it is also an ϵ -neighbourhood of $(B_\rho(1), d_\rho)$ and by corollary 6

$$|\|x_i - x_j\|_\infty - d_\rho(x_i, x_j)| \leq \epsilon.$$

Now let us take a continuous function from $B_\infty(1)$ to $\mathbb{R}$. Let $z_1, \dots, z_k$ and $\zeta_1, \dots, \zeta_l$ in the orbit of 0 by the $\mathbb{Z}^n$ action such that $\zeta_j + D \cap B_\infty(\rho) \neq \emptyset$ for $j = 1, \dots, l$ and

$$\bigcup_i z_i + D_f \subset B_\infty(\rho) \subset \bigcup_k \zeta_k + D_f$$

(where we took all z_i such that $z_i + D_f \subset B_\infty(\rho)$) then we get

$$\begin{aligned} \sum_i \inf_{\rho x \in z_i + D_f} f(x) \mu_\infty(D_f) &\leq \\ \int_{B_\infty(\rho)} f(x/\rho) d\mu_g(x) &\leq \sum_j \sup_{\rho x \in (\zeta_j + D_f) \cap B_\infty(\rho)} f(x) \mu_\infty(D_f) \end{aligned}$$

now dividing by ρ^n we find

$$\begin{aligned} \sum_i \inf_{x \in \frac{1}{\rho}(z_i + D_f)} f(x) \mu_\infty((1/\rho)D_f) &\leq \\ \int_{B_\infty(1)} f d\mu_\rho(x) &\leq \sum_j \sup_{x \in \frac{1}{\rho}(\zeta_j + D_f) \cap B_\infty(1)} f(x) \mu_\infty((1/\rho)D_f) \end{aligned}$$

The middle term is surrounded by two sums of Riemann, which converges to $\int_{B_\infty(1)} f d\mu_\infty$, thus it also converges. To conclude, notice that the net of characteristic function $\chi_{B_\rho(1)}$ converges simply to $\chi_{B_\infty(1)}$ inside of $B_\infty(1)$. $\square$

2.4 What shall we finally study ?

As we said we are now going to focus on the spectrum of the balls $B_g(\rho)$. As we allready mentioned we know that the eigenvalues are converging to zero with a $1/\rho^2$ speed. Hence we want to find a precise equivalent.

For this let introduce Δ_ρ the laplacian associated to the rescaled metrics $g_\rho = 1/\rho^2(\delta_\rho)^*g$, and for any function f from $B_g(\rho)$ to $\mathbb{R}$ lets associate a function f_ρ on $B_\rho(1)$ by $f_\rho(x) = f(\rho \cdot x)$. Then it is an easy calculation to see that for any $x \in B_\rho(1)$:

$$\rho^2(\Delta f)(\rho \cdot x) = (\Delta_\rho f_\rho)(x)$$

hence the eigenvalues of Δ_ρ on $B_\rho(1)$ are exactly the eigenvalues of Δ on $B_g(\rho)$ multiplied by ρ^2 and our problems becomes the study of the spectrum of the laplacian Δ_ρ on $B_\rho(1)$. In the light of what precedes we would like to show that there is some operator Δ_∞ acting on $B_\infty(1)$ such that, in some sense, the net of laplacian (Δ_ρ) converges towards Δ_∞ such that the spectra also converge to the spectrum of Δ_∞ . The following section aims at giving a precise meaning to this.

3 Convergence of spectral nets

This section adapts to our purpose some notion of convergences well known for a fixed Hilbert space.

3.1 Convergence on a net of Hilbert spaces

Let $(X_\alpha, d_\alpha, m_\alpha)_{\alpha \in \mathcal{A}}$, where $\mathcal{A}$ is a partially ordered set, be a net of compact measured metric spaces converging in the Gromov-Hausdorff measured topology to $(X_\infty, d_\infty, m_\infty)$. We will write $L_\alpha^2 = L^2(X_\alpha, m_\alpha)$ (resp. $L_\infty^2(X_\infty, m_\infty)$) for the square integrable function spaces. Their respective scalar product will be $\langle \cdot, \cdot \rangle_\alpha$ (resp. $\langle \cdot, \cdot \rangle_\infty$) and $\| \cdot \|_\alpha$ (resp. $\| \cdot \|_\infty$).

Furthermore we suppose that in every L_α^2 the continuous functions form a dense subset.

Definition 10. We say that a net $(u_\alpha)_{\alpha \in \mathcal{A}}$ of functions $u_\alpha \in L_\alpha^2$ strongly converges to $u \in L_\infty^2$ if there exists a net $(v_\beta)_{\beta \in \mathcal{B}} \subset C^0(X_\infty)$ converging to u in L_∞^2 such that

$$\lim_{\beta} \limsup_{\alpha} \|f_\alpha^* v_\beta - u_\alpha\|_\alpha = 0;$$

where (f_α) is the net of Hausdorff approximations. We will also talk of strong convergence in $\mathcal{L}^2$.

Definition 11. We say that a net $(u_\alpha)_{\alpha \in \mathcal{A}}$ of functions $u_\alpha \in L_\alpha^2$ weakly converges to $u \in L_\infty^2$ if and only if for every net $(v_\alpha)_{\alpha \in \mathcal{A}}$ strongly converging to $v \in L_\infty^2$ we have

$$\lim_{\alpha} \langle u_\alpha, v_\alpha \rangle_\alpha = \langle u, v \rangle_\infty \quad (3)$$

We will also talk of weak convergence in $\mathcal{L}^2$.

The following lemmas justifies those two definitions

Lemma 12. Let $(u_\alpha)_{\alpha \in \mathcal{A}}$ be a net of functions $u_\alpha \in L_\alpha^2$. If $(\|u_\alpha\|_\alpha)$ is uniformly bounded, then there exists a weakly converging subnet.

Proof. Let $(\phi_k)_{k \in \mathbb{N}}$ be a complete orthonormal basis of L_∞^2 . Using the density of continuous functions in L_∞^2 , for each k we can retrieve a net of continuous functions $(\varphi_{k,\beta})_{\beta \in \mathcal{B}}$ strongly converging to ϕ_k in L_∞^2 . Replacing by a subnet of $\mathcal{A}$ and $\mathcal{B}$ if necessarily, we can assume that the following limit exists

$$\lim_{\beta} \lim_{\alpha} \langle u_\alpha, f_\alpha^* \varphi_{1,\beta} \rangle_\alpha = a_1 \in \overline{\mathbb{R}}$$

and from the uniform bound hypothesis it follows that $a_1 \in \mathbb{R}$. Repeating this procedure we can assume that for every $k \in \mathbb{N}$ the following limit exists

$$\lim_{\beta} \lim_{\alpha} \langle u_{\alpha}, f_{\alpha}^* \varphi_{k,\beta} \rangle_{\alpha} = a_k \in \mathbb{R}.$$

Let us fix an integer N . For any $\epsilon > 0$ there is a $\beta_{\epsilon} \in \mathcal{B}$ such that

$$|\langle \varphi_{k,\beta}, \varphi_{l,\beta} \rangle_{\infty} - \delta_{kl}| < \epsilon$$

for any $\beta \geq \beta_{\epsilon}$ and $k, l = 1, \dots, N$. Moreover for any $\beta \geq \beta_{\epsilon}$ there is an $\alpha_{\epsilon,\beta} \in \mathcal{A}$ such that

$$|\langle f_{\alpha}^* \varphi_{k,\beta}, f_{\alpha}^* \varphi_{l,\beta} \rangle_{\alpha} - \delta_{kl}| < 2\epsilon$$

for any $\alpha \geq \alpha_{\epsilon,\beta}$ and $k, l = 1, \dots, N$. Let $L_{\alpha,\beta} = \text{Vect}\{f_{\alpha}^* \varphi_{k,\beta} \mid k = 1, \dots, N\}$ and $P_{\alpha,\beta} : L_{\alpha}^2 \rightarrow L_{\alpha,\beta}$ be the projection to the linear subspace $L_{\alpha,\beta} \subset L_{\alpha}^2$ we have

$$\left| \sum_{k=1}^N |\langle u_{\alpha}, f_{\alpha}^* \varphi_{k,\beta} \rangle_{\alpha}|^2 - \|P_{\alpha,\beta} u_{\alpha}\|_{\alpha}^2 \right| \leq \theta_N(\epsilon)$$

for every $\alpha \geq \alpha_{\epsilon,\beta}$ and $\beta \geq \beta_{\epsilon}$, where θ_N is a function depending only of N such that $\lim_{\epsilon \rightarrow 0} \theta_N(\epsilon) = 0$. This implies for every N

$$\begin{aligned} \sum_{k=1}^N |a_k|^2 &= \lim_{\beta} \lim_{\alpha} \sum_{k=1}^N |\langle u_{\alpha}, f_{\alpha}^* \varphi_{k,\beta} \rangle_{\alpha}|^2 = \lim_{\beta} \lim_{\alpha} \|P_{\alpha,\beta} u_{\alpha}\|_{\alpha}^2 \\ &\leq \limsup_{\alpha} \|u_{\alpha}\|_{\alpha}^2 < \infty \end{aligned}$$

thus

$$u = \sum_{k=1}^N a_k \phi_k \in L_{\infty}^2.$$

We shall prove that some subnet of $(u_{\alpha})_{\alpha}$ weakly converges to u . Take any $v \in L_{\infty}^2$ and set $b_k = \langle v, \phi_k \rangle_{\infty}$. By the properties of the strong convergence it is enough to show (3) for a well chosen net. Let $v_{\beta}^N = \sum_{k=1}^N b_k \varphi_{k,\beta}$. By construction $v_{\beta}^N \in \mathcal{C}^0$ and $\lim_{N \rightarrow \infty} \lim_{\beta} v_{\beta}^N = v$ strongly. We have

$$\lim_{\beta} \lim_{\alpha} \langle u_{\alpha}, f_{\alpha}^* v_{\beta}^N \rangle_{\alpha} = \lim_{\beta} \lim_{\alpha} \sum_{k=1}^N b_k \langle u_{\alpha}, f_{\alpha}^* \varphi_{k,\beta} \rangle_{\alpha} = \sum_{k=1}^N a_k b_k$$

which tends to $\langle u, v \rangle_{\infty}$ as $N \rightarrow \infty$. Thus, there exists a net of integers $(N_{\beta})_{\beta}$ tending to $+\infty$ such that $v_{\beta}^{N_{\beta}}$ strongly converges to v and

$$\lim_{\beta} \lim_{\alpha} \langle u_{\alpha}, f_{\alpha}^* v_{\beta}^{N_{\beta}} \rangle_{\alpha} = \langle u, v \rangle_{\infty}.$$

□

Lemma 13. *Let $(u_{\alpha})_{\alpha \in \mathcal{A}}$ be a weakly converging net to $u \in L_{\infty}^2$. Then*

$$\sup_{\alpha} \|u_{\alpha}\|_{\alpha} < \infty \quad \text{and} \quad \|u\|_{\infty} \leq \liminf_{\alpha} \|u_{\alpha}\|_{\alpha}.$$

Furthermore, the net strongly converges if and only if

$$\|u\|_{\infty} = \lim_{\alpha} \|u_{\alpha}\|_{\alpha}.$$

Proof. Let suppose that the (u_α) is weakly converging and $\sup_\alpha \|u_\alpha\|_\alpha = +\infty$. We can extract a sequence such that $\|u_{\alpha_k}\|_{\alpha_k} > k$. Setting

$$v_k = \frac{1}{k} \frac{u_{\alpha_k}}{\|u_{\alpha_k}\|_{\alpha_k}}$$

one has $\|v_k\|_{\alpha_k} = 1/k \rightarrow 0$ thus v_k strongly converges to 0, which implies

$$\langle u_{\alpha_k}, v_k \rangle_{\alpha_k} \rightarrow \langle u, 0 \rangle_\infty = 0$$

but we also have

$$\langle u_{\alpha_k}, v_k \rangle_{\alpha_k} = \frac{1}{k} \|u_{\alpha_k}\|_{\alpha_k} \geq 1$$

this is a contradiction and thus we obtain $\sup_\alpha \|u_\alpha\|_\alpha < \infty$.

Let $(w_\alpha)_\alpha$ be a strongly converging net to u , then

$$\begin{aligned} 0 &\leq \liminf_\alpha \|u_\alpha - w_\alpha\|_\alpha^2 \\ &= \liminf_\alpha (\|u_\alpha\|_\alpha^2 + \|w_\alpha\|_\alpha^2 - 2\langle u_\alpha, w_\alpha \rangle_\alpha) \\ &= \liminf_\alpha \|u_\alpha\|_\alpha^2 - \|u\|_\infty^2 \end{aligned}$$

The final claim comes from the properties of the strong convergence and the following equality

$$\|u_\alpha - w_\alpha\|_\alpha^2 = \|u_\alpha\|_\alpha^2 + \|w_\alpha\|_\alpha^2 - 2\langle u_\alpha, w_\alpha \rangle_\alpha.$$

□

3.2 Convergence of bounded operators

Let $\mathcal{L}(L_\alpha^2)$ be the set of linear bounded operators acting on L_α^2 and $\|\cdot\|_{\mathcal{L}_\alpha}$ their norm (for $\alpha \in \mathcal{A} \cup \infty$). Let $B_\infty \in \mathcal{L}(L_\infty^2)$ and $B_\alpha \in \mathcal{L}(L_\alpha^2)$ for every $\alpha \in \mathcal{A}$.

Theorem and Definition 14. Let $u, v \in L_\infty^2$ and $(u_\alpha)_{\alpha \in \mathcal{A}}, (v_\alpha)_{\alpha \in \mathcal{A}}$ two nets such that $u_\alpha, v_\alpha \in L_\alpha^2$. We say that the net of operators $(B_\alpha)_{\alpha \in \mathcal{A}}$ strongly (resp. weakly, compactly) converges to B if $B_\alpha u_\alpha \rightarrow Bu$ strongly (resp. weakly, strongly) for every net (u_α) strongly (resp. weakly, weakly) converging to $u \iff$

$$\lim_\alpha \langle B_\alpha u_\alpha, v_\alpha \rangle_\alpha = \langle Bu, v \rangle_\infty \quad (4)$$

for every $(u_\alpha), (v_\alpha), u$ and v such that $u_\alpha \rightarrow u$ strongly (resp. weakly, weakly) and $v_\alpha \rightarrow v$ weakly (resp. strongly, weakly).

Proof. The equivalence comes from the definition of the weak convergence and the fact that a net (u_α) strongly converges to u if and only if $\langle u_\alpha, v_\alpha \rangle_\alpha \rightarrow \langle u, v \rangle_\infty$ for every net $(v_\alpha)_\alpha$ weakly converging to $v \in L_\infty^2$. The "if" part is straightforward, for the "only if" we see that for every net (v_α) strongly converging to v we have $\langle u_\alpha, v_\alpha \rangle_\alpha \rightarrow \langle u, v \rangle_\infty$, which implies the weak convergence of the net (u_α) . Using now the hypothesis we get the convergence of the net $\|u_\alpha\|_\alpha$ and thus the strong convergence of (u_α) by lemma 13. □

Proposition 15. *Let (B_α) be a strongly converging net to B then*

$$\liminf_{\alpha} \|B_\alpha\|_{\mathcal{L}_\alpha} \geq \|B\|_{\mathcal{L}_\infty}$$

and if the convergence is compact then it is an equality and B is a compact operator as is its adjoint B^ .*

Proof. Let $\epsilon > 0$, there is $u \in \mathcal{L}_\infty^2$ such that $\|u\|_\infty = 1$ and $\|Bu\|_\infty > \|B\|_{\mathcal{L}_\infty} - \epsilon$. Take $(u_\alpha)_\alpha$ a net converging strongly to u . Then $\|u_\alpha\|_\alpha \rightarrow 1$, furthermore the strong convergence of (B_α) implies that $\|B_\alpha u_\alpha\|_\alpha \rightarrow \|Bu\|_\infty$ thus

$$\liminf_{\alpha} \|B_\alpha\|_{\mathcal{L}_\alpha} \geq \liminf_{\alpha} \frac{\|B_\alpha u_\alpha\|_\alpha}{\|u_\alpha\|_\alpha} = \|Bu\|_\infty > \|B\|_{\mathcal{L}_\infty} - \epsilon.$$

Let suppose now that the convergence is compact. Take a net (u_α) such that $\|u_\alpha\|_\alpha = 1$ and

$$\lim_{\alpha} |\|B_\alpha\|_{\mathcal{L}_\alpha} - \|B_\alpha u_\alpha\|_\alpha| = 0$$

extracting a subnet if necessary we can suppose that the net (u_α) weakly converges to u . By lemma 13 we have $\|u\|_\infty \leq 1$, furthermore the compact convergence implies the strong convergence of $B_\alpha u_\alpha$ to Bu thus

$$\|B\|_{\mathcal{L}_\infty} \geq \frac{\|Bu\|_\infty}{\|u\|_\infty} \geq \|Bu\|_\infty = \lim_{\alpha} \|B_\alpha u_\alpha\|_\alpha = \lim_{\alpha} \|B_\alpha\|_{\mathcal{L}_\alpha}$$

Now let us prove that in the latest case, B is compact. Let $(v_\beta)_{\beta \in \mathcal{B}}$ a net weakly converging to v in $\mathcal{L}_\infty^2$ then

$$\langle u, Bv_\beta \rangle_\infty = \langle B^* u, v_\beta \rangle_\infty \rightarrow \langle B^* u, v \rangle_\infty = \langle u, Bv \rangle_\infty$$

thus Bv_β weakly converges to Bv . For every β let $(u_{\alpha,\beta})$ be a strongly converging net such that $\lim_{\alpha} u_{\alpha,\beta} = v_\beta$. For every β the compact convergence of (B_α) implies the strong convergence of $B_\alpha u_{\alpha,\beta}$ to Bv_β . Now let us take a net of positive numbers such that $\lim_{\beta} \epsilon(\beta) = 0$, then there is $\alpha(\beta)$ such that for every $\alpha \geq \alpha(\beta)$ we have

$$|\|B_\alpha u_{\alpha,\beta}\|_\alpha - \|Bv_\beta\|_\infty| \leq \epsilon(\beta).$$

Set $w_\beta = u_{\alpha(\beta),\beta}$ then $\lim_{\beta} w_\beta = v$ weakly and by the compact convergence we obtain the strong convergence of $(B_{\alpha(\beta)} w_\beta)_\beta$ to Bv but

$$\lim_{\beta} |\|B_{\alpha(\beta)} w_\beta\|_{\alpha(\beta)} - \|Bv_\beta\|_\infty| = 0$$

which implies $\|Bv_\beta\|_\infty \rightarrow \|Bv\|_\infty$. We can conclude using lemma 13. $\square$

3.3 Convergence of spectral structures

Here we see L_α^2 as a Hilbert space. Then A_α and A will be self-adjoint operators, E_α and E their respective spectral measure and R_μ , R their resolvents for μ in the resolvent space. We want to study the links between the convergence of (A_α) , (E_α) and (R_μ^α) . The following theorem says that it is the same

Theorem 16. Let (A_α) and A be self-adjoint operators E_α, E their spectral measures and R_μ^α, R_μ their resolvents for μ in the resolvent space, then the following assertions are equivalent

1. $R_\mu^\alpha \rightarrow R_\mu$ strongly (resp. compactly) for μ outside the union of the spectra of A_α and A .
2. $\varphi(A_\alpha) \rightarrow \varphi(A)$ strongly (resp. compactly) for every continuous function, with compact support $\varphi : \mathbb{R} \rightarrow \mathbb{C}$
3. $\varphi_\alpha(A_\alpha) \rightarrow \varphi(A)$ strongly (resp. compactly) for every net $\{\varphi_\alpha : \mathbb{R} \rightarrow \mathbb{C}\}$ of continuous functions vanishing at infinity and uniformly converging to φ a continuous function vanishing at infinity.
4. $E_\alpha([\lambda, \mu]) \rightarrow E([\lambda, \mu])$ strongly (resp. compactly) for every pair of real numbers outside the spectrum of A .
5. $\langle E_\alpha u_\alpha, v_\alpha \rangle_\alpha \rightarrow \langle Eu, v \rangle_\infty$ vaguely for every net of vectors $(u_\alpha)_{\alpha \in \mathcal{A}}$ and $(v_\alpha)_{\alpha \in \mathcal{A}}$ such that $u_\alpha \rightarrow u$ strongly (resp. weakly) and $v_\alpha \rightarrow v$ weakly.

Let us recall that a quadratic form $\mathcal{Q}$ on a complex (resp. real) Hilbert space $\mathcal{H}$ comes from a sesquilinear (resp. bilinear) form, positive and symmetric $\mathcal{E} : D(\mathcal{E}) \times D(\mathcal{E}) \rightarrow \mathbb{C}$ (resp. $\mathbb{R}$) where $D(\mathcal{E}) \in \mathcal{H}$ is a linear subspace and $\mathcal{Q}(u) = \mathcal{E}(u, u)$. Notice that $\mathcal{E}_1(u, v) = \langle u, v \rangle_{\mathcal{H}} + \mathcal{E}(u, v)$ for every u and $v \in D(\mathcal{E})$ is also a sesquilinear (resp. bilinear), symmetric and positive form. Thus $((D(\mathcal{E}), \mathcal{E}_1))$ is a pre-Hilbert space. We say that $\mathcal{Q}$ is closed if and only if $((D(\mathcal{E}), \mathcal{E}_1))$ is a Hilbert space. In what follows, we will not distinguish $\mathcal{Q}$ and the functional $\mathcal{E}$ defined by $\mathcal{E}(u) = \mathcal{Q}(u)$ on $D(\mathcal{E})$ and $\mathcal{E}(u) = \infty$ on $\mathcal{H} \setminus D(\mathcal{E})$. In this context, $\mathcal{Q}$ is closed if and only if $\mathcal{E}$ is lower semi-continuous as a function $\mathcal{E} : \mathcal{H} \rightarrow \overline{\mathbb{R}}$.

Definition 17. Let $(\mathcal{E}_\alpha)$ be a net of closed quadratic forms, where $\mathcal{E}_\alpha$ is a closed quadratic form on L_α^2 for every $\alpha \in \mathcal{A}$. We will say that this net is asymptotically compact if and only if for every net $(v_\alpha)_{\alpha \in \mathcal{A}}$ such that

$$\limsup_{\alpha} \mathcal{E}_\alpha(v_\alpha) + \|v_\alpha\|_\alpha^2 < \infty$$

there is a strongly converging sub-net.

Now a **spectral structure** on a Hilbert space $\mathcal{H}$ over $\mathbb{C}$ (resp. $\mathbb{R}$) is a family

$$\Sigma = \{A, \mathcal{E}, E, (T_t), (R_\zeta)\}$$

where A is a self-adjoint operator seen as the infinitesimal generator of the densely defined quadratic form $\mathcal{E}$ (such that $D(\mathcal{E}) = D(\sqrt{A})$ and $\mathcal{E}(u, v) = \langle \sqrt{A}u, \sqrt{A}v \rangle_{\mathcal{H}}$ for every u and v in $D(\mathcal{E})$), E is its spectral measure, $(T_t)_{t \geq 0}$ is a one parameter semi-group of strongly continuous contractions ($T_t = e^{-tA}$, $t \geq 0$) and R_ζ is a strongly continuous resolvent ($R_\zeta = (\zeta - A)^{-1}$ for $\zeta \in \rho(A)$, where $\rho(A)$ is the resolvent set of A). In what follows we will study a family of spectral structures Σ_α on L_α^2 , thus we will have

$$\Sigma_\alpha = \{A_\alpha, \mathcal{E}_\alpha, E_\alpha, (T_t^\alpha), (R_\zeta^\alpha)\}$$

Definition 18. Let $(\Sigma_\alpha)_{\alpha \in \mathcal{A}}$ be a net with Σ_α a spectral structure on L_α^2 and Σ a spectral structure on L_∞^2 , we will say that the net $(\Sigma_\alpha)_\alpha$ strongly (resp.

compactly) converges to Σ if and only if one of the conditions of theorem 16 is satisfied.

Proposition 19. *Let $(\Sigma_\alpha)_{\alpha \in \mathcal{A}}$ of spectral structures strongly converging to Σ then for any net $(v_\alpha)_\alpha$ weakly converging to v we have*

$$\mathcal{E}(v) \leq \liminf_\alpha \mathcal{E}_\alpha(v_\alpha)$$

Furthermore, if the net $(\Sigma_\alpha)_{\alpha \in \mathcal{A}}$ converges compactly, then the net of quadratic forms $(\mathcal{E}_\alpha)_\alpha$ is asymptotically compact.

Proof. Assume that the net of resolvents (R_λ^α) is strongly convergent and write

$$a_\alpha^\lambda(u, v) = -\lambda \langle u - \lambda R_\lambda^\alpha u, v \rangle_\alpha$$

(the Deny-Yosida approximation of bilinear form associated to $\mathcal{E}_\alpha$), then the net $(a_\alpha^\lambda(u, u))$ converges to $\mathcal{E}_\alpha(u)$ increasing when $\lambda \rightarrow -\infty$ (see Mosco [Mos94] 1.(i)). From the assumption it is easy to see that for (u_α) and (v_α) converging strongly to u and weakly to v respectively

$$\lim_\alpha a_\alpha^\lambda(u_\alpha, v_\alpha) = -\lambda \langle u - \lambda R_\lambda u, v \rangle_\infty = a^\lambda(u, v)$$

we recall that (see Dal Maso [Mas93] proposition 12.12)

$$a^\lambda(u, u) \geq a^\lambda(v, v) + 2\lambda \langle v - \lambda R_\lambda v, u - v \rangle_\infty$$

hence for any net v_α weakly converging to u and w_α a strongly converging net to u we have

$$\mathcal{E}_\alpha(v_\alpha) \geq a_\alpha^\lambda(v_\alpha, v_\alpha) \geq a_\alpha^\lambda(w_\alpha, w_\alpha) + 2\lambda \langle w_\alpha - \lambda R_\lambda^\alpha w_\alpha, v_\alpha - w_\alpha \rangle$$

thus $\liminf_\alpha \mathcal{E}_\alpha(v_\alpha) \geq a^\lambda(u, u)$ for any $\lambda < 0$, now taking $\lambda \rightarrow -\infty$ we can conclude that $\liminf_\alpha \mathcal{E}_\alpha(v_\alpha) \geq \mathcal{E}(u)$.

Now assume that (Σ_α) compactly converges and let $(u_\alpha)_{\alpha \in \mathcal{A}}$ be a net such that

$$\sup_\alpha (\mathcal{E}_\alpha(u_\alpha) + \|u_\alpha\|_\alpha^2) \leq M < \infty .$$

Taking a sub-net if necessary we can suppose that $(u_\alpha)_\alpha$ weakly converges to u . Let $\rho > 0$ be out of A_∞ 's spectrum. As

$$\int_{[\rho, \infty[} d\langle E_\alpha u_\alpha, u_\alpha \rangle_\alpha \leq \frac{1}{\rho} \int_{[\rho, \infty[} \lambda d\langle E_\alpha(\lambda) u_\alpha, u_\alpha \rangle_\alpha \leq \frac{\mathcal{E}_\alpha(u_\alpha)}{\rho} \leq \frac{M}{\rho}$$

we have

$$\|u_\alpha\|_\alpha^2 \leq \int_{[0, \rho]} d\langle E_\alpha u_\alpha, u_\alpha \rangle_\alpha + \frac{M}{\rho}$$

and the compact convergence implies $\lim_\alpha \int_{[0, \rho]} d\langle E_\alpha u_\alpha, u_\alpha \rangle_\alpha = \int_{[0, \rho]} d\langle Eu, u \rangle_\infty$ hence

$$\limsup_\alpha \|u_\alpha\|_\alpha^2 \leq \int_{[0, \rho]} d\langle Eu, u \rangle_\infty + \frac{M}{\rho} \leq \|u\|_\infty^2 + \frac{M}{\rho}$$

and taking $\rho \rightarrow \infty$ we get

$$\limsup_\alpha \|u_\alpha\|_\alpha^2 \leq \|u\|_\infty^2$$

finally we deduce the strongly convergence of the net (u_α) using lemma 13. $\square$

The main reason we introduced all these convergences is the following theorem, the proof of which we postpone to avoid drowning the reader in too many technical details.

Theorem 20. *Let $\Sigma_\alpha \rightarrow \Sigma$ compactly and suppose that all resolvents R_ζ^α are compact. Let λ_k (resp. λ_k^α) be the k^{th} eigenvalue of A (resp. A_α) with multiplicity. We take $\lambda_k = +\infty$ if $k > \dim L_\infty^2 + 1$ when $\dim L_\infty^2 < \infty$ and $\lambda_k^\alpha = +\infty$ if $k > \dim L_\alpha^2 + 1$ when $\dim L_\alpha^2 < \infty$. Then for every k*

$$\lim_{\alpha} \lambda_k^\alpha = \lambda_k$$

Furthermore let $\{\varphi_k^\alpha \mid k = 1, \dots, \dim L_\alpha^2\}$ be an orthonormal bases of L_α^2 such that φ_k^α is an eigenfunction of A_α for λ_k^α . Then there is a sub-net such that for all $k \leq \dim L_\infty^2$ the net $(\varphi_k^\alpha)_\alpha$ strongly converges to the eigenfunction φ_k of A for the eigenvalue λ_k , and such that the family $\{\varphi_k \mid k = 1, \dots, \dim L_\infty^2\}$ is an orthonormal basis of L_∞^2 .

4 Proof of Theorem 1

4.1 Homogenisation of the Laplacian

In this section we are going to build the operator Δ_∞ of theorem 1. we remind the reader that D_f is a fundamental domain, we then begin by taking χ^i as the unique periodic solution (up to an additive constant) of

$$\Delta \chi^i = \Delta x_i \text{ on } D_f$$

The operator Δ_∞ is then defined by

$$\Delta_\infty f = -\frac{1}{\text{Vol}(g)} \left(\int_{D_f} g^{ij} - g^{ik} \frac{\partial \chi^j}{\partial y_k} d\mu_g \right) \frac{\partial^2 f}{\partial x_i \partial x_j} \quad (5)$$

Now let us write $\eta_j(x) = \chi^j(x) - x_j$ the induced harmonic function and

$$q^{ij} = \frac{1}{\text{Vol}(g)} \left(\int_{D_f} g^{ij} - g^{ik} \frac{\partial \chi^j}{\partial y_k} d\mu_g \right)$$

we can notice that the $d\eta_i$ are harmonic 1-forms on the torus. It is not difficult now to show that

Proposition 21. *Let $\langle \cdot, \cdot \rangle_2$ be the scalar product induced on 1-forms by the Riemannian metric g . Then*

$$q^{ij} = \frac{1}{\text{Vol}(g)} \langle d\eta_i, d\eta_j \rangle_2 = q^{ji}$$

thus Δ_∞ is an elliptic operator.

In fact we can say more, (q^{ij}) induces a scalar product on harmonic 1-forms (whose norm will be written $\|\cdot\|_2$) and then to $H^1(\mathbb{T}, \mathbb{R})$. Indeed, as mentioned earlier, we can see the $(d\eta_i)$ as 1-forms over the torus. Being a free family they can be seen as a basis of $H^1(\mathbb{T}, \mathbb{R})$ (Hodge's theorem). Thus by duality this yields also a scalar product (q_{ij}) over $H_1(\mathbb{T}, \mathbb{R})$ (whose induced norm will be written $\|\cdot\|_2^*$). The question naturally arising is to know the link between this norm and the stable norm. To see this we have to go back on $H^1(\mathbb{T}, \mathbb{R})$. Indeed the stable norm is the dual of the norm obtained by quotient of the sup norm on 1-forms (see Pansu [Pan99] lemma 17), which we write $\|\cdot\|_\infty^*$, and the norm $\|\cdot\|_2$ comes from the normalised L^2 norm. Thus mixing the Hölder inequality and the Hodge-de Rham theorem we get :

Proposition 22. *For every 1-form α we have*

$$\|\alpha\|_2 \leq \|\alpha\|_\infty^*$$

thus by duality, for every $\gamma \in H_1(\mathbb{T}, \mathbb{R})$ we have

$$\|\gamma\|_\infty \leq \|\gamma\|_2^*$$

in other words the unit ball of $\|\cdot\|_2^$ is included in $B_\infty(1)$.*

To finish this section, let us remark that the manifold $H_1(\mathbb{T}, \mathbb{R})/H_1(\mathbb{T}, \mathbb{Z})$ with the flat metric induced by $\|\cdot\|_2^*$ is usually called the Jacobi manifold or the Albanese torus of $(\mathbb{T}, g)$.

4.2 Asymptotic compactness

Let us now define the various functional spaces involved. For $\rho \in \overline{\mathbb{R}}$, $L^2(B_\rho(1), d\mu_\rho)$ will be the space of square integrable functions over the ball $B_\rho(1)$, which is a Hilbert space with the scalar product

$$(u, v)_\rho = \int_{B_\rho(1)} uv \, d\mu_\rho$$

whose norm will be $|\cdot|_\rho$. $H_{\rho,0}^1(B_\rho(1))$ will be the closure of $C^\infty(B_\rho(1))$ functions with compact support, in $H_\rho^1(B_\rho(1))$ for the norm $\|\cdot\|_\rho$ defined by

$$\|v\|_\rho^2 = |v|_\rho^2 + \sum_{i=1}^n \left| \frac{\partial v}{\partial x_i} \right|_\rho^2$$

and with

$$H_\rho^1(B_\rho(1)) = \left\{ v \left| v, \frac{\partial v}{\partial x_1}, \dots, \frac{\partial v}{\partial x_n} \in L^2(B_\rho(1), d\mu_\rho) \right. \right\}$$

For all that follows, V_ρ will be a closed sub-space such that

$$H_{\rho,0}^1(B_\rho(1)) \subset V_\rho \subset H_\rho^1(B_\rho(1))$$

Thus we can define a spectral structure on L_ρ^2 by expanding the Laplacian defined on V_ρ on L_ρ^2 . If $V_\rho = H_{\rho,0}^1(B_\rho(1))$ we deal with the Dirichlet problem, and if $V_\rho = H_\rho^1(B_\rho(1))$ we then deal with the Neumann problem. We then put the following norm on V_ρ :

$$\|v\|_{\rho,0}^2 = |v|_\rho^2 + (v, \Delta_\rho v)_\rho$$

we then have

Lemma 23. *Let $(u_\rho)_\rho$ be a net with $u_\rho \in V_\rho$ for every ρ , if there is a constant C such that for all $\rho > 0$ we have*

$$\|u_\rho\|_{\rho,0} \leq C$$

then there is a strongly converging sub-net in $\mathcal{L}^2$.

Proof. Let $B = \cup_\rho B_\rho(1)$ we are going to show that the strong convergence in $L^2(B, \mu_\infty)$ implies the strong convergence in $\mathcal{L}^2$. Then the compact embedding of $H_\infty^1(B)$ in $L^2(B, \mu_\infty)$ will conclude the proof.

Let us first notice that the periodicity gives the existence of two constant α and β such that

$$\alpha|v|_\infty \leq |v|_\rho \leq \beta|v|_\infty .$$

Let us start by taking a net (u_ρ) strongly converging in $L^2(B, \mu_\infty)$ to u_∞ we also assume $u_\rho \in V_\rho$ for every ρ , because it is all we need. Now let $c_p \in C_0^\infty(B_\infty(1))$ be a sequence of functions strongly converging to u_∞ and take p large enough for the support of u_p to be in $B_\rho(1)$. We have

$$|c_p - u_\rho|_\rho \leq \beta|c_p - u_\infty|_\infty + \beta|u_\infty - u_\rho|_\infty$$

now let $\varepsilon > 0$ then for p large enough $\beta|c_p - u_\infty|_\infty \leq \varepsilon$. We fix p large enough and take ρ large enough for the second term to converge to 0.

Now to conclude observe that from the assumptions the net (u_ρ) is bounded in $H_\infty^1(B)$, hence using the compact embedding of $H_\infty^1(B)$ in $L^2(B, \mu_\infty)$ we can extract a strongly converging net in $L^2(B, \mu_\infty)$ and by what we just did in $\mathcal{L}^2$. $\square$

4.3 Compact convergence of the resolvents

Let $\lambda > 0$ and G_λ^ρ be the operator from L_ρ^2 to $V_\rho \subset L_\rho^2$ such that

$$a_\lambda^\rho(G_\lambda^\rho f, \phi) = (f, \phi)_\rho \quad \forall \phi \in V_\rho. \quad (6)$$

where

$$a_\lambda^\rho(u, v) = \int_{B_\rho(1)} g_\rho^{ij} \partial_i u \cdot \partial_j v \, d\mu_\rho + \lambda(u, v)_\rho$$

We want to show that the net of operators (G_λ^ρ) converges compactly to G_λ the operator corresponding to the homogenised problem:

$$a_\lambda^\infty(G_\lambda f, \phi) = (f, \phi)_\infty \quad \forall \phi \in V_\infty \quad (7)$$

with $(f, \phi)_\infty = \int_{B_\infty(1)} f \phi \, d\mu_\infty$ and

$$a_\lambda^\infty(u, v) = \int_{B_\infty(1)} q^{ij} \partial_i u \partial_j v \, d\mu_\infty + \lambda(u, v)_\infty$$

in other word we want to show the following theorem

Theorem 24. *For every $\lambda < 0$, the net of resolvents $(R_\lambda^\rho)_\rho$ of the Laplacian (Δ_ρ) converges compactly to R_λ^∞ , the resolvent of Δ_∞ from the homogenised problem. Thus the net (Σ_ρ) compactly converges to Σ_∞ .*

Proof. This comes from the fact that $R_\lambda^\rho = -G_{-\lambda}^\rho$ and $R_\lambda^\infty = -G_{-\lambda}$.

First step :

Let f_ρ be a weakly convergent net to f in $\mathcal{L}^2$, thus from 13 this net is uniformly bounded in $\mathcal{L}^2$ and in V_ρ' , the dual space of V_ρ .

Let $f_\rho \in V_\rho$ then by (6) we have :

$$\alpha \|G_\lambda^\rho f_\rho\|_{\rho,0}^2 \leq (f_\rho, G_\lambda^\rho f_\rho)_\rho \leq K \|f_\rho\|_{V_\rho'} \|G_\lambda^\rho f_\rho\|_{\rho,0}$$

thus

$$\|G_\lambda^\rho f_\rho\|_{\rho,0} \leq C \|f_\rho\|_{V_\rho'}$$

the net $(G_\lambda^\rho f_\rho)$ being uniformly bounded for the norms $\|\cdot\|_{\rho,0}$, using lemma 12 there is a subnet strongly converging in $\mathcal{L}^2$. i.e.

$$u_\rho = G_\lambda^\rho f_\rho \rightarrow u_\lambda^* \text{ strongly in } \mathcal{L}^2 \quad (8)$$

Furthermore $P_\rho = (g_\rho^{ij}) \nabla G_\lambda^\rho f_\rho$ is also bounded in $\mathcal{L}^2$ thus there is a subnet of the net P_ρ weakly converging in $\mathcal{L}^2$ to $P_\lambda^* \in L_\infty^2$. For any $\phi_\infty \in L_\infty^2$ let ϕ_ρ be a strongly converging net to ϕ_∞ in $\mathcal{L}^2$ then

$$\begin{aligned} \int_{B_\rho(1)} P_\rho \cdot \nabla \phi_\rho \, d\mu_\rho + \lambda (G_\lambda^\rho f_\rho, \phi_\rho)_\rho &= (f_\rho, \phi_\rho)_\rho \rightarrow \\ \int_{B_\infty(1)} P_\lambda^* \cdot \nabla \phi_\infty \, d\mu_\infty + \lambda (u_\lambda^*, \phi_\infty)_\infty &= (f, \phi_\infty)_\infty. \end{aligned} \quad (9)$$

Thus it is enough to show that $P_\lambda^* = (q^{ij}) \nabla u_\lambda^*$ on $B_\infty(1)$ because it induces $u_\lambda^* = G_\lambda f$.

Second step :

We first take $\chi^k(y)$ (see 4.1) such that $\mathcal{M}(\chi^k) = 0$ and we define

$$w_\rho(x) = x_k - \frac{1}{\rho} \chi^k(\rho x) \quad (10)$$

for every $k = 1, \dots, d_1$. Then

$$w_\rho \rightarrow x_k \text{ strongly in } \mathcal{L}^2. \quad (11)$$

an by construction of χ^k (see 4.1) we have

$$-\partial_i (F(g_\rho, X) g_\rho^{ij} \partial_j w_\rho) = 0 \text{ on } B_\rho(1). \quad (12)$$

We multiply this equation by a test function $\phi \in V_\rho$ and after an integration we get

$$\int_{B_\rho(1)} g_\rho^{ij} \partial_j w_\rho \partial_i \phi \, d\mu_\rho = 0 \quad (13)$$

Let $\varphi \in C_0^\infty(B_\infty(1))$ (notice that for ρ large enough the support of φ will be in $B_\rho(1)$) and $\phi = \varphi w_\rho$ which we put into the equation (6) and into the equation (13) we put $\phi = \varphi u_\rho$ (see (8)), and then we subtract the results

$$\begin{aligned} \int_{B_\rho(1)} g_\rho^{ij} (\partial_j u_\rho \partial_i \varphi w_\rho - \partial_j w_\rho \partial_i \varphi u_\rho) \, d\mu_\rho \\ = \int_{B_\rho(1)} f_\rho w_\rho \varphi \, d\mu_\rho - \lambda \int_{B_\rho(1)} \varphi u_\rho w_\rho \, d\mu_\rho \end{aligned} \quad (14)$$

Now let $\rho \rightarrow \infty$ in (14), all terms converge because they are product of one strongly converging net and one weakly converging net in $\mathcal{L}^2$. More precisely

- P_ρ defined $P_{\rho,i} = g_\rho^{ij} \partial_j u_\rho$ weakly converges to P_λ^* in $\mathcal{L}^2$ following (9);
- $\partial_i \varphi w_\rho$ strongly converges to $\partial_i \varphi x_k$ in $\mathcal{L}^2$ from (11)
- $g_\rho^{ij} \partial_i w_\rho$ is D_f/ρ -periodic and weakly converges in $\mathcal{L}^2$ towards its mean value

$$q^{jk} = \mathcal{M} \left(g^{ij}(y) \left(\delta_{ik} - \partial_i \chi^k(y) \right) \right)$$

- $\partial_j \varphi u_\rho$ strongly converges to $\partial_j \varphi u_\lambda^*$ by (8), because φ has compact support.
- Now for the right side, w_ρ strongly converges as u_ρ does and f_ρ weakly converges to f .

To summarise (14) converges to (we write $P_{\lambda,i}^*$ the coordinates of P_λ^*)

$$\int_{B_\infty(1)} (P_{\lambda,j}^* x_k - q^{jk} u_\lambda^*) \partial_j \varphi \, d\mu_\infty = \int_{B_\infty(1)} f x_k \varphi \, d\mu_\infty - \lambda \int_{B_\infty(1)} \varphi u_\lambda^* x_k \, d\mu_\infty \quad (15)$$

furthermore if we put into equation (9), $\phi_\infty = \varphi x_k$ it gives

$$\int_{B_\infty(1)} f x_k \varphi \, d\mu_\infty - \lambda \int_{B_\infty(1)} \varphi u_\lambda^* x_k \, d\mu_\infty = \int_{B_\infty(1)} P_{\lambda,j}^* \partial_j (\varphi x_k) \, d\mu_\infty \quad (16)$$

and by mixing (15) and (16) we get for every $\varphi \in \mathcal{C}_c^\infty(B_\infty(1))$ the following equality :

$$\int_{B_\infty(1)} (P_{\lambda,j}^* x_k - q^{jk} u_\lambda^*) \partial_j \varphi \, d\mu_\infty = \int_{B_\infty(1)} P_{\lambda,j}^* \partial_j (\varphi x_k) \, d\mu_\infty$$

which in terms of distribution can be translated into :

$$-\sum_{j=1}^{d_1} \partial_j (P_{\lambda,j}^* x_k - q^{jk} u_\lambda^*) = -\sum_{j=1}^{d_1} \partial_j P_j^* x_k \iff P_{\lambda,k}^* = \sum_{j=1}^{d_1} q^{jk} \partial_j u_\lambda^*$$

which allow us to conclude that $u_\lambda^* = G_\lambda f$. $\square$

It is now easy to finish the proof of theorem 1, it comes from theorem 24 and theorem 20.

5 Γ -convergence of quadratic forms

5.1 Γ and Mosco-convergence of quadratic forms

We are now going to give a definition of Γ -convergence adapted to our problem.

Definition 25. (Γ -convergence). We say that a net $\{F_\alpha : L_\alpha^2 \rightarrow \overline{\mathbb{R}}\}_{\alpha \in \mathcal{A}}$ of functions Γ -converges to $F : L_\infty^2 \rightarrow \overline{\mathbb{R}}$ if and only if the following assertions are satisfied :

(F1) For any net $(u_\alpha)_{\alpha \in \mathcal{A}} \in L_\alpha^2$ strongly converging to $u \in L_\infty^2$ in $\mathcal{L}^2$ we have

$$F(u) \leq \liminf_\alpha F_\alpha(u_\alpha);$$

(F2) For every $u \in L_\infty^2$ there is a net $(u_\alpha)_{\alpha \in \mathcal{A}} \in L_\alpha^2$ strongly converging to u in $\mathcal{L}^2$ such that

$$F(u) = \lim_\alpha F_\alpha(u_\alpha).$$

Remark. This is slightly different from the definition 5, which is the usual one. By taking F_α infinite outside of L_α^2 in $\mathcal{L}^2$ we get back (in some way) the usual definition (see the introduction of [Mas93]).

Let us summarise some properties satisfied by this convergence.

Lemma 26.

- (a) Let $\{F_\alpha : L_\alpha^2 \rightarrow \overline{\mathbb{R}}\}_{\alpha \in \mathcal{A}}$ be a net of functions Γ -converging to a function $F : L_\infty^2 \rightarrow \overline{\mathbb{R}}$, then F is lower semi-continuous.
- (b) Let $(\mathcal{E}_\alpha)_{\alpha \in \mathcal{A}}$ be a net of quadratic forms $\mathcal{E}_\alpha$ on L_α^2 Γ -converging to a function $F : L_\infty^2 \rightarrow \overline{\mathbb{R}}$, then F can be identified with a quadratic form on L_∞^2 .

There is also the following result, concerning compactness

Theorem 27. From every net $(\mathcal{E}_\alpha)_{\alpha \in \mathcal{A}}$ of quadratic forms $\mathcal{E}_\alpha$ on L_α^2 we can extract a Γ -converging subnet, whose limit is a quadratic form on L_∞^2 .

Remark. This theorem is true for a wider variety of functions, with some restrictions on $\{L_\alpha^2\}_{\alpha \in \mathcal{A}}$. Of course the limit in that case is not always a quadratic form. Here it is lemma 26 which gives information on the limit.

Definition 28. (Mosco topology). We say that a net $(\mathcal{E}_\alpha)_{\alpha \in \mathcal{A}}$ of quadratic forms $\mathcal{E}_\alpha$ on L_α^2 Mosco-converges to the quadratic form $\mathcal{E}$ on L_∞^2 if condition (F2) of definition 25 and (F1') are satisfied:

(F1') For any $(u_\alpha)_{\alpha \in \mathcal{A}}$, $u_\alpha \in L_\alpha^2$ weakly converging net to $u \in L_\infty^2$ in $\mathcal{L}^2$ we have

$$\mathcal{E}(u) \leq \liminf_{\alpha} \mathcal{E}_\alpha(u_\alpha)$$

The induced topology is called the Mosco topology.

It is obvious that the Mosco-convergence induces the Γ -convergence, thus this topology is stronger. Let us now define one last convergence :

Definition 29. (Compact Γ -convergence). We say that a net $(\mathcal{E}_\alpha)_{\alpha \in \mathcal{A}}$ Γ -converges compactly to $\mathcal{E}$ if $\mathcal{E}_\alpha \rightarrow \mathcal{E}$ in the Mosco topology and if $(\mathcal{E}_\alpha)_{\alpha \in \mathcal{A}}$ is asymptotically compact.

Let us precise how the Mosco and the Γ topologies are linked :

Lemma 30. Let us suppose $(\mathcal{E}_\alpha)_{\alpha \in \mathcal{A}}$ asymptotically compact then $(\mathcal{E}_\alpha)_{\alpha \in \mathcal{A}}$ Γ -converges to $\mathcal{E}$ if and only if $(\mathcal{E}_\alpha)_{\alpha \in \mathcal{A}}$ Mosco-converges to $\mathcal{E}$.

Proof. We just need to show that the Γ -convergence implies the condition (F1') from definition 28. We proceed *ad absurdum* and suppose that there is a weakly converging net (u_α) such that $\liminf_{\alpha} \mathcal{E}_\alpha(u_\alpha) < \mathcal{E}(u)$. Taking a subnet if necessarily we can suppose $\lim_{\alpha} \mathcal{E}_\alpha(u_\alpha) < \mathcal{E}(u)$ thus we also have $\limsup_{\alpha} \mathcal{E}_\alpha(u_\alpha) + \|u_\alpha\|_{\alpha}^2 < +\infty$. The asymptotic compactness is obviously inherited by a subnet thus we can extract a strongly converging subnet $u_{\alpha(\beta)}$. The Γ -convergence being also inherited by a subnet of $\mathcal{E}_\alpha$ we finally get

$$\lim_{\beta} \mathcal{E}_\alpha(u_{\alpha(\beta)}) = \lim_{\beta} \mathcal{E}_{\alpha(\beta)}(u_{\alpha(\beta)}) \geq \mathcal{E}(u)$$

which is absurd. □

5.2 Γ -convergence and spectral structures

The following theorem explains how are related the convergence of spectral structures and the Mosco-convergence.

Theorem 31. *Let (Σ_α) be a net of spectral structures on (L_α^2) and Σ a spectral structure on L_∞^2 then $\Sigma_\alpha \longrightarrow \Sigma$ strongly (resp. compactly) if and only if $\mathcal{E}_\alpha$ Mosco-converges (resp. Γ -converges compactly) to $\mathcal{E}$.*

Proof. We are going to prove the equivalence between the strong (resp. compact) convergence of resolvents and the Mosco-convergence (resp. Γ -convergence compact) of the energies.

Let us begin by assuming the Mosco-convergence of the net $(\mathcal{E}_\alpha)$. We need to show that for every $z \in L_\infty^2$ and any net (z_α) strongly converging to z the net $u_\alpha = -R_\lambda^\alpha z_\alpha$ strongly converges to $u = -R_\lambda z$. First let us notice that the vector u is the unique minimiser of

$$v \mapsto \mathcal{E}(v) - \lambda \|v\|_\infty^2 - 2\langle z, v \rangle_\infty$$

we can characterise the same way u_α for every α .

As an operator of L_α^2 , R_λ^α is bounded by $-\lambda^{-1}$. Thus the net (u_α) is bounded and we can extract a weakly converging sub-net, still written (u_α) , with limit $\tilde{u}$. Now from condition (F2) for every $v \in L_\infty^2$ there is a net strongly converging to it such that $\lim_\alpha \mathcal{E}_\alpha(v_\alpha) = \mathcal{E}(v)$. But for every α

$$\mathcal{E}_\alpha(u_\alpha) - \lambda \|u_\alpha\|_\alpha^2 - 2\langle z_\alpha, u_\alpha \rangle_\alpha \leq \mathcal{E}_\alpha(v_\alpha) - \lambda \|v_\alpha\|_\alpha^2 - 2\langle z_\alpha, v_\alpha \rangle_\alpha \quad (17)$$

thus taking the limit in $\alpha \in \mathcal{A}$ we get thanks to condition (F1') of definition 28 and the fact that for any weakly convergent net $\|\tilde{u}\|_\infty \leq \liminf_\alpha \|u_\alpha\|_\alpha$ (remember that $\lambda < 0$)

$$\mathcal{E}(\tilde{u}) - \lambda \|\tilde{u}\|_\infty^2 - 2\langle z, \tilde{u} \rangle_\infty \leq \mathcal{E}(v) - \lambda \|v\|_\infty^2 - 2\langle z, v \rangle_\infty$$

which implies $\tilde{u} = -R_\lambda z$. Due to u 's unicity, we conclude that (u_α) weakly converges to u . Let us prove that $\|u_\alpha\|_\alpha$ converges to $\|u\|_\infty$. In that aim take a strongly convergent net v_α to v such that $\lim_\alpha \mathcal{E}_\alpha(v_\alpha) = \mathcal{E}(u)$, and take a new look at inequality (17) :

$$\mathcal{E}_\alpha(u_\alpha) - \lambda \|u_\alpha + z_\alpha/\lambda\|_\alpha^2 \leq \mathcal{E}_\alpha(v_\alpha) - \lambda \|v_\alpha + z_\alpha/\lambda\|_\alpha^2$$

using (F1') once again we find

$$\mathcal{E}(v) - \lambda \limsup_\alpha \|u_\alpha + z_\alpha/\lambda\|_\alpha^2 \leq \mathcal{E}(v) - \lambda \|u + z/\lambda\|_\infty^2$$

thus $\|u_\alpha + z_\alpha/\lambda\|_\alpha^2 \rightarrow \|u + z/\lambda\|_\infty^2$ which implies the strong convergence of $(u_\alpha + z_\alpha/\lambda)_\alpha$ and the strong convergence of (z_α) induces the strong convergence of (u_α) .

We shall now study the compact Γ -convergence. Let us take a weakly convergent net w_α to w and let $u_\alpha = -R_\lambda^\alpha w_\alpha$, then the net u_α is still bounded. Swapping z_α with w_α in (17) we get that $\limsup_\alpha \mathcal{E}_\alpha(u_\alpha)$ is bounded, and thanks to the asymptotic compactness we can extract a strongly convergent sub-net with $\tilde{u}$ its limit. Putting this in (17), with $z_\alpha = w_\alpha$ where (w_α) a strongly converging net to w we get

$$\mathcal{E}(\tilde{u}) - \lambda \|\tilde{u}\|^2 - 2\langle w, \tilde{u} \rangle \leq \mathcal{E}(v) - \lambda \|v\|^2 - 2\langle w, v \rangle$$

thus $\tilde{u} = -R_\lambda w$. Once again, thanks to unicity, we conclude that $R_\lambda^\alpha w_\alpha$ strongly converges to $R_\lambda w$.

Reciprocally assume that for every $\lambda < 0$ the net R_λ^α strongly converges to R_λ . In what follows (u_α) will be a strong convergent net to u .

Condition (F1') : already done, see proposition 19.

Condition (F2) : Extract a sub-net $\lambda_\alpha \rightarrow -\infty$ such that

$$\mathcal{E}(u, u) \geq \lim_{\lambda} \lim_{\alpha} a_{\alpha}^{\lambda}(u_{\alpha}, u_{\alpha}) \geq \lim_{\alpha} a_{\alpha}^{\lambda_{\alpha}}(u_{\alpha}, u_{\alpha})$$

take $w_{\alpha} = \lambda_{\alpha} R_{\lambda_{\alpha}}^{\alpha} u_{\alpha}$ for every α and notice that

$$\begin{aligned} a_{\alpha}^{\lambda}(u_{\alpha}, u_{\alpha}) &= -\lambda \langle u_{\alpha} - \lambda R_{\lambda}^{\alpha} u_{\alpha}, u_{\alpha} \rangle_{\alpha} - \lambda \langle u_{\alpha} - \lambda R_{\lambda}^{\alpha} u_{\alpha}, -\lambda R_{\lambda}^{\alpha} u_{\alpha} \rangle_{\alpha} \\ &\quad + \lambda \langle u_{\alpha} - \lambda R_{\lambda}^{\alpha} u_{\alpha}, -\lambda R_{\lambda}^{\alpha} u_{\alpha} \rangle_{\alpha} \\ &= -\lambda \|u_{\alpha} - \lambda R_{\lambda}^{\alpha} u_{\alpha}\|^2 + \lambda^2 \langle u_{\alpha} - \lambda R_{\lambda}^{\alpha} u_{\alpha}, -R_{\lambda}^{\alpha} u_{\alpha} \rangle_{\alpha} \\ &= -\lambda \|u_{\alpha} - \lambda R_{\lambda}^{\alpha} u_{\alpha}\|^2 + \mathcal{E}_{\alpha}(\lambda R_{\lambda}^{\alpha} u_{\alpha}) \end{aligned}$$

indeed if a_{α} is the bilinear form corresponding to $\mathcal{E}_{\alpha}$ then $R_{\lambda}^{\alpha} u_{\alpha}$ can be seen as the sole element such that

$$a_{\alpha}(-R_{\lambda}^{\alpha} u_{\alpha}, v_{\alpha}) - \lambda \langle -R_{\lambda}^{\alpha} u_{\alpha}, v_{\alpha} \rangle_{\alpha} = \langle u_{\alpha}, v_{\alpha} \rangle_{\alpha}, \quad \forall v_{\alpha} \in D(\mathcal{E}_{\alpha})$$

hence

$$a_{\alpha}^{\lambda_{\alpha}}(u_{\alpha}, u_{\alpha}) = \mathcal{E}_{\alpha}(w_{\alpha}, w_{\alpha}) - \lambda_{\alpha} \|u_{\alpha} - w_{\alpha}\|^2$$

which implies $w_{\alpha} \rightarrow u$ strongly in $\mathcal{L}^2$ and

$$\mathcal{E}(u, u) \geq \limsup_{\alpha \rightarrow +\infty} \mathcal{E}_{\alpha}(w_{\alpha}, w_{\alpha})$$

For the compact convergence case it suffices to prove the asymptotic compactness, but it has already been done in the proof of proposition 19. $\square$

6 Proof of Theorem 2

The convergence of the eigenvalue is given by theorem 1. Hence it remains to bound the asymptotic λ_1 (i.e. the limit) and characterise the equality. The proof we propose consists in finding an upper bound of $\lambda_1(B_g(\rho))$ for every ρ using a function depending of the distance from the centre of the ball. We then use the simple convergence of the distances (d_{ρ}) to the stable norm as seen in section 2.2 and the measure part of theorem 9.

Proof. Let f be a continuous function from $\mathbb{R}$ to $\mathbb{R}$ and define

$$\begin{aligned} f_{\rho} : B_{\rho}(1) &\longrightarrow \mathbb{R} \\ x &\longmapsto f(d_{\rho}(0, x)) \end{aligned}$$

and $f_{\infty}(x) = f(\|x\|_{\infty})$ on $B_{\infty}(1)$. We want to show that (remember that $\delta_{\rho}(x) = \rho x$)

$$\int f_{\rho} \cdot \chi_{B_g(\rho)} \circ \delta_{\rho} d\mu_{\rho} \xrightarrow{\rho \rightarrow \infty} \int_{B_{\infty}(1)} f_{\infty} d\mu_{\infty} \quad (18)$$

To obtain this we are going to cut the difference in three pieces i.e. :

$$\left| \int f_\rho \cdot \chi_{B_g(\rho)} \circ \delta_\rho d\mu_\rho - \int_{B_\infty(1)} f_\infty d\mu_\infty \right| \leq \left| \int f_\rho \cdot (\chi_{B_g(\rho)} \circ \delta_\rho - \chi_{B_\infty(1)}) d\mu_\rho \right| \quad (19)$$

$$+ \left| \int_{B_\infty(1)} (f_\rho - f_\infty) d\mu_\rho \right| \quad (20)$$

$$+ \left| \int_{B_\infty(1)} f_\infty d\mu_\rho - \int_{B_\infty(1)} f_\infty d\mu_\infty \right| \quad (21)$$

Now it suffices to notice that

- 1) the part (19) goes to 0 because inside we have the product of $\chi_{B_g(\rho)} \circ \delta_\rho - \chi_{B_\infty(1)}$, which is easily seen to simply converge to 0 thanks to 8, with bounded terms compactly supported.
- 2) Same reason but here it is $f_\rho - f_\infty$ which simply converges to 0,
- 3) Finally the convergence to 0 of (21) is due once again to the measure part of theorem 9.

As a conclusion we have (18). Injecting now f_ρ into the Raleigh's quotient we get :

$$\rho^2 \lambda_g(B_g(\rho)) \leq \frac{\int ((f')_\rho)^2 \cdot \chi_{B_g(\rho)} \circ \delta_\rho d\mu_\rho}{\int (f_\rho)^2 \cdot \chi_{B_g(\rho)} \circ \delta_\rho d\mu_\rho}$$

we apply (18) to obtain

$$\limsup_{\rho \rightarrow \infty} \rho^2 \lambda_g(B_g(\rho)) \leq \frac{\int_{B_\infty(1)} ((f')_\infty)^2 d\mu_\infty}{\int_{B_\infty(1)} f_\infty^2 d\mu_\infty}$$

and now taking for f the right function (i.e. the solution of the differential equation $f'' + \frac{n-1}{x} f'(x) + \lambda_{e,n} f = 0$) we can conclude.

Let us now study the equality case. Take again the function f which gives the eigenfunction of the Euclidean Laplacian on the Euclidean unit ball (i.e. the solution of $f'' + \frac{n-1}{x} f'(x) + \lambda_{e,n} f = 0$) and normalise it. The Γ -convergence theory allows to say, taking $\mathcal{E}_\rho$ and $\mathcal{E}_\infty$ as the energies of Δ_ρ and Δ_∞ on the balls $B_\rho(1)$ and $B_\infty(1)$ respectively for the adapted measures and thanks to proposition 19 and 1

$$\mathcal{E}_\infty(f_\infty) \leq \liminf_{\rho \rightarrow \infty} \mathcal{E}_\rho(f_\rho) \leq \limsup_{\rho \rightarrow \infty} \mathcal{E}_\rho(f_\rho) \leq \lambda_{e,n} \quad (22)$$

Now from the equality assumption we have

$$\lambda_{e,n} \leq \mathcal{E}_\infty(f_\infty), \quad (23)$$

thus (22) and (23) imply equality which in turn imply that f_∞ is an eigenfunction for the first eigenvalue. Hence f_∞ is smooth (at least in a neighbourhood of zero).

Now from the study of Bessel's function (see. bow, §103 à §105)) we see that taking $p = (n - 2)/2$ we have $f(x) = x^{-p} J_p(\sqrt{\lambda_e} x)$ with J_p an analytic function defined by (see F. Bowman [Bow58] §84)

$$J_p(x) = \frac{x^p}{2^p \Gamma(p+1)} \left(1 - \frac{x^2}{2 \cdot 2n+2} + \frac{x^4}{2 \cdot 4 \cdot 2n+2 \cdot 2n+4} + \dots \right)$$

thus f has the following shape

$$f(x) = \frac{\lambda_e^p}{2^p \Gamma(p+1)} \left(1 - \frac{x^2 \lambda_e}{2 \cdot 2n+2} + \frac{x^4 \lambda_e^2}{2 \cdot 4 \cdot 2n+2 \cdot 2n+4} + \dots \right)$$

in other words f has the following asymptotic expansion $1 + \alpha_1 x^2 + \alpha_2 x^4 + \dots$ (up to a multiplicative constant), now notice that the function $1 + \alpha_1 x + \alpha_2 x^2 + \dots$ admits an inverse $g \in C^\infty$ in a neighbourhood of zero, which implies that $g \circ f_\infty(x) = cst \cdot \|x\|_\infty^2$ is C^2 in a neighbourhood of zero, thus the stable norm comes from a scalar products, which means that it is Euclidean.

In fact we have some more informations. Indeed in order for f_∞ to be an eigenfunction, the norme of the differential of the stable norm with respect to the albanese metric (the scalar product giving the laplacian Δ_∞) must be almost everywhere equal to one (a simple computation using the fact that the stable norm is euclidean and the Cauchy-Schwartz inequality). Which implies that the unit ball of the albanese metric must be inside the unit ball of the stable norm. Now the maximum principle and the monotony with respect to inclusion of the eigenvalues implies that equality holds if and only if the stable norm and the Albanese metric coincides. The stable norm being the Albanese metric we can now use the theorem 32 to conclude. $\square$

Theorem 32. *Let $(\mathbb{T}^n, g)$ be a torus, its stable norm coincides with the Albanese metric if and only if the torus is flat.*

Proof. Let us take a base $\eta_1, \dots, \eta_n$ of Harmonic one forms, any function α and any 2-form β . We shall write $(\cdot, \cdot)_g$ pointwise scalar product induced by g on forms ($\|\cdot\|_g$ the associated norm) and $\langle \cdot, \cdot \rangle_g$ the integral scalar product normalized by the volume. Then by Hodge's theorem

$$\|\eta_i\|_\infty^2 = \inf_{\alpha, \beta} \sup_{x \in \mathbb{T}^n} \|\eta_i\|_g^2 + \|d\alpha\|_g^2 + \|\delta\beta\|_g^2 \geq \sup_{x \in \mathbb{T}^n} \|\eta_i\|_g^2$$

and

$$\langle \eta_i, \eta_i \rangle_g = \frac{1}{\text{Vol}_g(\mathbb{T}^n)} \int_{\mathbb{T}^n} \|\eta_i\|_g^2 d\text{vol}_g \leq \sup_{x \in \mathbb{T}^n} \|\eta_i\|_g^2$$

the case of equality implies that $\langle \eta_i, \eta_i \rangle_g = (\eta_i, \eta_i)_g(x)$ for all $x \in \mathbb{T}^n$. Now it suffices to see that the metric g can be written in the following way :

$$\sum_{i,j} \lambda_{ij} \eta_i \circ \eta_j = g$$

where $\eta_i \circ \eta_j = 1/2(\eta_i \otimes \eta_j + \eta_j \otimes \eta_i)$ and $\Lambda = (\lambda_{ij})$ is the matrice such that $\Lambda^{-1} = (\langle \eta_i, \eta_j \rangle_g)$. Now taking local f_i such that $df_i = \eta_i$ then the function $F(x) = (f_1(x), \dots, f_n(x))$ is an isometry between a open set of $\mathbb{T}^n$ and an euclidean space, thus the torus is flat. $\square$

7 Related topics

In that section we come back to the asymptotic volume, proving in the meantime a generalised Faber-Krahn inequality. Then we explain what can be deduced from our work for the heat kernel and how it is related to other's work. We finally state how theorem 1 passes to graded nilmanifolds.

7.1 Asymptotic volume of tori

7.1.1 Generalised Faber-Krahn inequality

We need some more definitions

Definition 33. For a rectifiable submanifold N of $\mathbb{R}^n$ (we can think of it of finite adapted Hausdorff measure) we will write $I(N)$ the associated integral current. For an integral current C , $M(C)$ will be its mass as defined par H. Federer (see [Fed69] for example).

Definition 34. Let $\mathbb{R}^n$, with the norm $\|\cdot\|$ ($\|\cdot\|_*$ will be the dual norm), we define

$$\lambda_1(\Omega, \|\cdot\|) = \inf_f \frac{\int_{\Omega} \|df\|_*^2 d\mu}{\int_{\Omega} f^2 d\mu}$$

where μ is the Lebesgue measure on $\mathbb{R}^n$, and the infimum is taken over all lipschitz functions vanishing on the border.

The following lemma holds,

Lemma 35. (Faber-Krahn inequality for norms). Let D be a domain of $\mathbb{R}^n$, with the norm $\|\cdot\|$ and a measure μ invariant by translation. Let D^* be the norm's ball with same measure as D , then

$$\lambda_1(D^*, \|\cdot\|) \leq \lambda_1(D, \|\cdot\|) \quad (24)$$

the equality case implying that D is a norm's ball.

Proof. We need two ingredients for this proof. The first is an isoperimetric inequality, which is given by a result of Brunn (see a proof by M. Gromov in [MS86]). The second is a co-area formula, which can be found in Federer [Fed69] p. 438.

More specifically, let us write $G_t = \{x \mid |f(x)| = t\}$ then on one side we have

$$\int_{\Omega} h\alpha \wedge df = \int_0^{\sup f} \int_{G_t} h\alpha|_{G_t} dt = \int_0^{\sup f} I_{|f|=t}(h\alpha) dt$$

and on the other

$$\int_{\Omega} \|df\|_*^2 d\mu = \int_0^{\sup f} M(I_{|f|=t}) dt \quad (25)$$

(see P. Pansu [Pan99]) where $d\mu$ is the translation invariant volume form on $\mathbb{R}^n$ such that the norm's ball of radius one has measure 1.

Take $\alpha = \frac{1}{|df|^2} * df$ where $*$ is the Hodge operator on differential forms over $\mathbb{R}^n$ then we get

$$\int_{\Omega} h d\mu = \int_0^{\sup f} \int_{G_t} h \alpha|_{G_t} dt$$

Take now a look at the same equality on $\Omega_t = \{x \mid |f(x)| > t\}$ i.e. :

$$\int_{\Omega_t} h d\mu = \int_t^{\sup f} \int_{G_t} h \alpha|_{G_t} dt = \int_t^{\sup f} I_{|f|=t}(h\alpha) dt \quad (26)$$

differentiating each member of equality (26) we get almost everywhere the following equality

$$\int_{G_t} h \alpha|_{G_t} = I_{|f|=t}(h\alpha) \quad (27)$$

taking into account (27) and (25) we obtain

$$\int_{G_t} \|df\|_* \alpha|_{G_t} = M(I_{|f|=t}) \quad (28)$$

applying the Cauchy-Schwartz inequality to the left side of (28) and make the appropriate identification thanks to (27) we finally have

$$\frac{M(I_{|f|=t})^2}{I_{|f|=t}(\alpha)} \leq I_{|f|=t}((\|df\|_*)^2 \alpha) \quad (29)$$

The function f^* associated to f by symmetrisation is lipschitz, thus she satisfies a similar co-area formula. Hence we have for almost all t

$$I_{|f|=t}(\alpha) = -\frac{d}{dt} \text{Vol}(\Omega_t) = -\frac{d}{dt} \text{Vol}(\Omega_t^*) = I_{|f^*|=t}(\alpha^*) \quad (30)$$

now using the Brunn's isoperimetric inequality (see [MS86]) we have

$$M(I_{|f^*|=t}) \leq M(I_{|f|=t}) \quad (31)$$

injecting (30) and (31) in (29) and noticing that $\|df^*\|_*$ is constant on $\{|f|=t\}$, which implies that the equivalent of (29) for f^* is an equality we get (for almost all t)

$$I_{|f^*|=t}((\|df^*\|_*)^2 \alpha^*) = \frac{M(I_{|f^*|=t})^2}{I_{|f^*|=t}(\alpha^*)} \leq \frac{M(I_{|f|=t})^2}{I_{|f|=t}(\alpha)} \leq I_{|f|=t}((\|df\|_*)^2 \alpha) \quad (32)$$

Now we sum the extremal terms of (32) to obtain the wanted inequality :

$$\int_{\Omega^*} (\|df^*\|_*)^2 dv \leq \int_{\Omega} (\|df\|_*)^2 dv$$

which allows us to conclude the proof because

$$\int_{\Omega^*} (f^*)^2 dv = \int_{\Omega} (f)^2 dv$$

For the equality case, it suffices to see that it implies the equality case in the Brunn's isoperimetric inequality to conclude. $\square$

Let us notice that this lemma immediately implies

Corollary. *Let D_1 the unit ball of the norm $\|\cdot\|$ then*

$$\lambda_1(D_1) = \lambda_{e,n}$$

thus

$$\lambda_{e,n} \left(\frac{\mu(D_1)}{\mu(D)} \right)^{\frac{2}{n}} \leq \lambda_1(D, \|\cdot\|)$$

where μ is a Haar measure on $\mathbb{R}^n$.

Proof. The symmetrisation from the previous theorem shows that the minimum of the Rayleigh's quotient is obtained with functions depending on the distance from the centre of the ball. Hence we are lead to the same calculations as in the Euclidean case. $\square$

7.1.2 Lower bound for the asymptotic volume

We are now going to apply the generalised Faber-Krahn inequality to λ_∞ . In that aim in mind let us notice that $\lambda_\infty(B_\infty(1)) = \lambda_1(B_\infty(1), \|\cdot\|_2^*)$ with the dual norm of $\|\cdot\|_2^*$ defined by

$$\|\xi\|_2 = \sum_{ij} q^{ij} \xi_i \xi_j$$

and let us write B_{Al} the unit ball of $\|\cdot\|_2^*$. We now can apply inequality of lemma 35 and more precisely its corollary (32)

$$\lambda_\infty(B_\infty(1)) \geq \left(\frac{\mu(B_{Al})}{\mu(B_\infty(1))} \right)^{2/n} \lambda_{e,n}$$

where μ is any Haar measure. Now applying theorem we get

$$\left(\frac{\mu(B_{Al})}{\mu(B_\infty(1))} \right)^{2/n} \lambda_{e,n} \leq \lambda_{e,n} \quad (33)$$

We finally get the following proposition taking in (33) the Haar measure such that the measure of $B_\infty(1)$ is the asymptotic volume (i.e. the measure μ_∞) and transforming the other term in order to make the Albanese's Torus's volume appear.

Proposition 3. *Let $(\mathbb{T}^n, g)$ be a Riemannian Torus, $B_g(\rho)$ the geodesic balls of radius ρ centred on a fixed point and $\text{Vol}_g(B_g(\rho))$ their Riemannian volume induced on the universal cover, writing*

$$\text{Asvol}(g) = \lim_{\rho \rightarrow \infty} \frac{\text{Vol}_g(B_g(\rho))}{\rho^n}$$

then

$$1. \text{ Asvol}(g) \geq \frac{\text{Vol}_g(\mathbb{T}^n)}{\text{Vol}_{Al}(\mathbb{T}^n)} \omega_n$$

2. *In case of equality, the torus is flat. Where ω_n is the unit Euclidean ball's Euclidean volume.*

Proof. There remains the equality case to be proved, which can be obtained using either the equality case of the Faber-Krahn inequality, which says that $B_\infty(1)$ is an ellipsoid either the equality case of theorem 2 and then we conclude by using theorem 32. $\square$

We still have two remarks concerning this proposition, the first one is included in the following corollary.

Corollary. *For $n = 2$ we have*

1. $\text{Asvol}(g) \geq \pi = \omega_2$
2. *In case of equality, the torus is flat.*

In other words we obtain the theorem of D. Burago et S. Ivanov on the asymptotic volume of tori in the 2 dimensional case (see [BI95]). The second remark is that we can not do better this way. See [Ver01] part three for more details.

7.2 Long time asymptotics of the heat kernel

Let $(\mathbb{T}^n, g)$ be a torus and $(\mathbb{R}^n, \tilde{g})$ its universal cover with the lifted metric. We remind the reader that $g_\rho = (1/\rho^2)\delta_\rho^*\tilde{g}$ are the rescaled metrics and Δ_ρ their Laplacian, here it will be on $\mathbb{R}^n$.

We are going to study from the homogenisation point of view the long time asymptotic behaviour of the heat kernel i.e. we are interested in the behaviour as t goes to infinity of a solution $u(t, x)$ of the following problem

$$\begin{cases} \frac{\partial u}{\partial t} + \Delta u = 0 & \text{in }]0, +\infty[\times \mathbb{R}^n \\ u(0, x) = u_0(x) \end{cases} \quad (34)$$

For a probabilistic insight one could see M. Kotani et T. Sunada [KS00].

Let us introduce the rescaled functions

$$u_\rho(t, x) = \rho^n u(\rho^2 t, \delta_\rho x), \quad \rho > 0$$

it is the straightforward that (see. 2.4) u is a solution of (34) if and only if u_ρ is a solution of

$$\begin{cases} \frac{\partial u_\rho}{\partial t} + \Delta_\rho u_\rho = 0 & \text{in }]0, +\infty[\times \mathbb{R}^n \\ u_\rho(0, x) = \rho^n u_0(\delta_\rho x) \end{cases} \quad (35)$$

hence studying $u(t, \cdot)$ as t goes to infinity is the same as studying $u_\rho(1, \cdot)$ as $\rho \rightarrow \infty$. In other words we are once again lead to the study of the spectral structures (Δ_ρ) on $\mathbb{R}^n$. We have

Theorem 36. *The net of resolvents (R_λ^ρ) weakly converges to the resolvent (R_λ^∞) of Δ_∞ in $L^2(\mathbb{R}^n)$.*

Remark. The proof is the same as 24. In fact in that case we would rather talk of G -convergence. We now can apply the theorems from the chapter III of [ZKON79] more precisely theorems 4 and 6.

Theorem 37. ([ZKON79] page 136). *The fundamental solution $k(t, x, y)$ of (34) has the following asymptotic expansion*

$$k(t, x, y) = k_\infty(t, x, y) + t^{-\frac{n}{2}} \theta(t, x, y)$$

where $k_\infty(t, x, y)$ is fundamental solution of

$$\frac{\partial u_\infty}{\partial t} + \Delta_\infty u_\infty = 0 \text{ in }]0, +\infty[\times \mathbb{R}^n \quad (36)$$

and $\theta(t, x, y) \rightarrow 0$ uniformly as $t \rightarrow \infty$ on $|x|^2 + |y|^2 \leq at$, for any fixed constant $a > 0$.

Remark. This is slightly weaker than theorem 1 of M. Kotani and T. Sunada in [KS00].

Theorem 38. ([ZKON79] page 138). *Let $u_0 \in L^1(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$. Then $u(t, x)$ the solution of (34) has the following asymptotic expansion :*

$$u(t, x) = c_0 (4\pi t)^{-\frac{n}{2}} \int_{\mathbb{R}^n} u_0(y) dy + t^{-\frac{n}{2}} \theta(t, x)$$

where $\theta(t, x)$ converges uniformly to 0 for $|x| < R$ where R is a positive constant and c_0 is the determinant of the matrix associated to Δ_∞ .

That last claim can be precised by

Theorem 39. (Duro, Zuazua [DZ00]). *Let $u_0 \in L^1(\mathbb{R}^n)$. The sole solution of (34) satisfies for every $p \in [1, +\infty[$:*

$$t^{n/2(1-1/p)} \|u(t) - u_\infty(t)\|_p \rightarrow 0, \text{ as } t \rightarrow +\infty \quad (37)$$

where u_∞ is the unique solution of the homogenised problem (36). For $n = 1$ and $n = 2$ (37) is also true for $p = \infty$.

7.3 The macroscopical sound of graded nilmanifolds

In this part we want to emphasise the fact that theorem 1 is still true for graded nilmanifolds, at least for the Dirichlet case, but it involves some sub-riemannian geometry. We just give the statement the details are to be found in [Ver01] chapter two.

Theorem 40. *Let (M^n, g) be graded nilmanifold, $B_g(\rho)$ the induced Riemannian ball of radius ρ on its universal cover and $\lambda_i(B_g(\rho))$ the i^{th} eigenvalue of the Laplacian on $B_g(\rho)$ for the Dirichlet problem.*

Then there exists an hypoelliptic operator Δ_∞ (the Kohn Laplacian of a left invariant metric), whose i^{th} eigenvalue for the Dirichlet problem on the stable ball is λ_i^∞ and such that

$$\lim_{\rho \rightarrow \infty} \rho^2 \lambda_i(B_g(\rho)) = \lambda_i^\infty$$

where the stable ball is the metric ball given by the Carnot-Caratheodory distance found in [Pan82] and arising from the stable norm.

8 Proof of theorem 20

Let (Σ_α) be a net of spectral structures and let us focus on the spectra. For a fixed operator $\sigma(\cdot)$ will be its spectrum. Let us begin with the case of strong convergence.

Proposition 41. *If $\Sigma_\alpha \rightarrow \Sigma$ strongly, then for any $\lambda \in \sigma(A)$ there is $\lambda_\alpha \in \sigma(A_\alpha)$ such that the net (λ_α) converges to λ , this is written*

$$\sigma(A) \subset \lim_{\alpha} \sigma(A_\alpha)$$

Proof. Let $\lambda \in \sigma(A)$ and $\varepsilon > 0$ and take $\zeta = \lambda + i\varepsilon$ then :

$$\|R_\zeta^\alpha\|_{\mathcal{L}_\alpha} = \frac{1}{\inf_{\rho \in \sigma(A_\alpha)} |\zeta - \rho|} \quad \text{and} \quad \|R_\zeta\|_{\mathcal{L}_\infty} = \frac{1}{\inf_{\rho \in \sigma(A)} |\zeta - \rho|} = \frac{1}{\varepsilon}.$$

From the assumption, the net of resolvents strongly converges hence by 15

$$\limsup_{\alpha} \inf_{\rho \in \sigma(A_\alpha)} |\zeta - \rho| \leq \varepsilon$$

and as it is true for any ε , we can conclude. □

Lemma 42. *For any reals a, b out of the spectra of A such that $-\infty \leq a < b \leq +\infty$ then*

$$a \leq \frac{\mathcal{E}(u)}{\|u\|_\infty^2} \leq b \quad \text{pour tout } u \in E([a, b])L_\infty^2 \setminus \{0\}.$$

(where $E([a, b]) = E([a, +\infty[)$ si $b = +\infty$).

Proof. Let $a < b$ two reals out of the spectra of A and

$$u \in E([a, b])L_\infty^2 \setminus \{0\}.$$

then

$$\int_{[a, b]} dEu = E([a, b])u = u = \int_{\mathbb{R}} dEu$$

thus $\langle Eu, u \rangle = 0$ on $\mathbb{R} \setminus]a, b]$. Now if $u \in D(A)$,

$$\mathcal{E}(u) = \langle Au, u \rangle = \int_{\mathbb{R}} \lambda d\langle E(\lambda)u, u \rangle = \int_{[a, b]} \lambda d\langle E(\lambda)u, u \rangle$$

and the last term satisfies

$$a\|u\|_\infty^2 = a \int_{[a, b]} d\langle E(\lambda)u, u \rangle \leq \int_{[a, b]} \lambda d\langle E(\lambda)u, u \rangle \leq b \int_{[a, b]} d\langle E(\lambda)u, u \rangle = b\|u\|_\infty^2$$

□

For any Borel set $I \subset \mathbb{R}$ we write $n(I) = \dim E(I)L_\infty^2$ and $n_\alpha(I) = \dim E_\alpha(I)L_\alpha^2$.

Proposition 43. *Let $a < b$ two reals out of the point spectrum of A . if $\Sigma_\alpha \rightarrow \Sigma$ strongly then*

$$\liminf_{\alpha} n_{\alpha}([a, b]) \geq n([a, b])$$

and in particular,

$$\liminf_{\alpha} \dim L_{\alpha}^2 \geq \dim L_{\infty}^2$$

Proof. Lets take an orthonormal base $\{\varphi_k \mid k = 1, \dots, n([a, b])\}$ of $E([a, b])L_{\infty}^2$. Let $n \in \mathbb{N}$ a fixed number if $n([a, b]) = \infty$ else $n = n([a, b])$. Then there are nets $\varphi_k^{\alpha} \in L_{\alpha}^2$ for $k = 1, \dots, n$ such that $\lim_{\alpha} \varphi_k^{\alpha} = \varphi_k$. As $E_{\alpha}([a, b]) \rightarrow E([a, b])$ strongly, taking $\psi_k^{\alpha} = E_{\alpha}([a, b])\varphi_k^{\alpha}$ we get

$$\lim_{\alpha} \psi_k^{\alpha} = E([a, b])\varphi_k = \varphi_k$$

hence

$$\lim_{\alpha} \langle \psi_i^{\alpha}, \psi_j^{\alpha} \rangle_{\alpha} = \langle \varphi_i, \varphi_j \rangle = \delta_{ij}$$

from which we deduce that $(\psi_k^{\alpha})_{k=1, \dots, n}$ is a free family for α large enough and

$$\lim_{\alpha} n_{\alpha}([a, b]) \geq n.$$

which proves the first assertion. For the second it comes from the fact that $n([a, b])$ converges to $\dim L_{\infty}^2$ as $a \rightarrow -\infty$ and $b \rightarrow +\infty$. $\square$

Let us now have a look at the compact convergence case :

Theorem 44. *If $\Sigma_{\alpha} \rightarrow \Sigma$ compactly converges, then for any a, b out of the point spectrum of A such that $a < b$ for α large enough we have $n_{\alpha}([a, b]) = n([a, b])$. In particular the limit of the sets $\sigma(A_{\alpha})$ coincides with $\sigma(A)$*

Proof. The compact convergence implies that the operators R_{ζ} , T_t and $E([\lambda, \mu])$ are compact (see 15) Thus the spectrum of A is discrete and $n([a, b]) < \infty$ if $a < b < \infty$. Let $(0 \leq) \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$ be the spectrum of A , where

$$\begin{cases} n = 0 & \text{if the spectre is empty ,} \\ n \in \mathbb{N} & \text{if the spectre is finite and} \\ n = \infty & \text{if the spectre is a sequence converging to infinity.} \end{cases}$$

Step 1 : Fix ε_0 and let $\Lambda_1^{\alpha} = E([-\infty, \lambda_1 + \varepsilon_0])L_{\alpha}^2$ and $\Lambda_1 = L_{\infty}^2$, where $\lambda_1 = \lambda_1 + \varepsilon_0 = \infty$ if $n = 0$. Let

$$\mu_1 = \liminf_{\alpha} \inf \{ \mathcal{E}_{\alpha}(u) \mid \|u\|_{\alpha} = 1, u \in \Lambda_1^{\alpha} \}$$

lemma 42 allows us to say that $\lim_{\alpha} n_{\alpha}([-\infty, \mu]) = 0$ for any $\mu \in]-\infty, \mu_1[$. Applying proposition 43 we get $n([-\infty, \mu]) = 0$, in other words for any $\mu \leq \mu_1$ then $\mu \leq \lambda_1$ thus $\mu_1 \leq \lambda_1$. Hence if $\mu_1 = +\infty$, $n = 0$ and $L_{\alpha}^2 = 0$ for α large enough and the theorem is proved in that case.

Suppose that $\mu_1 < +\infty$. For α large enough we can find unit vectors $\varphi_1^{\alpha} \in \Lambda_1^{\alpha}$ such that $\liminf_{\alpha} \mathcal{E}_{\alpha}(\varphi_1^{\alpha}) = \mu_1$. From the asymptotic compactness of $\mathcal{E}_{\alpha}$ we can extract a sub-net $(\varphi_1^{\alpha})_{\alpha \in \mathcal{A}}$ such that $\varphi_1 = \lim_{\alpha} \varphi_1^{\alpha}$ strongly and thanks to

definition 19 $\mathcal{E}(\varphi_1) \leq \mu_1$. The strong convergence induces the convergence of the norms hence $\|\varphi_1\| = 1$ and

$$\lambda_1 = \inf\{\mathcal{E}(u) \mid \|u\| = 1, u \in \Lambda_1\} \leq \mathcal{E}(\varphi_1) \leq \mu_1 < +\infty.$$

As a consequence $n \geq 1$, $\lambda_1 = \mu_1 = \mathcal{E}(\varphi_1)$ and φ_1 is eigenvector of A for λ_1 .

Furthermore let us notice that as $E_\alpha([\lambda_1 - \epsilon, \lambda_1 + \epsilon]) \rightarrow E([\lambda_1 - \epsilon, \lambda_1 + \epsilon])$ strongly for any $\epsilon > 0$ fixed and $E([\lambda_1 - \epsilon, \lambda_1 + \epsilon]) \rightarrow E(\{\lambda_1\})$ strongly when $\epsilon \rightarrow 0$ there is a net of positives numbers $\epsilon_1^\alpha \rightarrow 0$ such that $E_\alpha([\lambda_1 - \epsilon_1^\alpha, \lambda_1 + \epsilon_1^\alpha]) \rightarrow E(\{\lambda_1\})$ strongly. From this we obtain a net

$$\psi_1^\alpha = E_\alpha([\lambda_1 - \epsilon_1^\alpha, \lambda_1 + \epsilon_1^\alpha])\varphi_1^\alpha \rightarrow E(\{\lambda_1\})\varphi_1 = \varphi_1.$$

Step 2 : Let $\Lambda_2^\alpha = E([- \infty, \lambda_2 + \epsilon_0])L_\alpha^2 \cap \langle \varphi_1^\alpha \rangle^\perp$, $\Lambda_2 = \langle \varphi_1 \rangle^\perp$ and

$$\mu_2 = \liminf_\alpha \inf\{\mathcal{E}_\alpha(u) \mid \|u\|_\alpha = 1, u \in \Lambda_2^\alpha\}.$$

Again lemma 42 allows us to say that $\lim_\alpha n_\alpha([- \infty, \mu]) = 0$ for any $\mu \in ([\mu_1, \mu_2[)$ and proposition 43 that $\mu_2 \leq \lambda_2$. Hence if $\mu_2 = +\infty$, we have $n = 1$ and $L_\alpha^2 = \langle \psi_1^\alpha \rangle$ for α large enough. Assume $\mu_2 < \infty$. Take the unitary vectors $\varphi_2^\alpha \in \Lambda_2^\alpha$ such that $\liminf_\alpha \mathcal{E}_\alpha(\varphi_2^\alpha) = \mu_2$. Then the same discussion as step 1 gives $n \geq 2$, $\lambda_2 = \mu_2$ and the strong convergence of a sub-net of (φ_2^α) to φ_2 an eigenvector of A for the eigenvalue λ_2 . We also find a net $\epsilon_2^\alpha \rightarrow 0$ such that $\psi_2^\alpha = E_\alpha([\lambda_2 - \epsilon_2^\alpha, \lambda_2 + \epsilon_2^\alpha])L_\alpha^2 \rightarrow \varphi_2$. Now let us notice that for any $\epsilon > 0$ there is $\alpha_\epsilon \in \mathcal{A}$ such that for all $\alpha \geq \alpha_\epsilon$ we have

$$(1) \psi_i^\alpha \in E_\alpha([\lambda_i - \epsilon, \lambda_i + \epsilon])L_\alpha^2 \text{ for } i = 1, 2 ;$$

$$(2) \text{ if } \lambda_1 + 2\epsilon < \lambda_2 \text{ then}$$

$$E_\alpha([\lambda_1 - \epsilon, \lambda_1 + \epsilon])L_\alpha^2 = \langle \psi_1^\alpha \rangle \quad \text{and} \quad E_\alpha([\lambda_1 + \epsilon, \lambda_2 - \epsilon])L_\alpha^2 = 0.$$

Step 3 : We repeat this procedure. Setting

$$\Lambda_k^\alpha = E([- \infty, \lambda_k + \epsilon_0])L_\alpha^2 \cap \langle \psi_1^\alpha, \dots, \psi_{k-1}^\alpha \rangle^\perp$$

we have

$$\lambda_k = \mu_k = \liminf_\alpha \inf\{\mathcal{E}_\alpha(u) \mid \|u\|_\alpha = 1, u \in \Lambda_k^\alpha\}$$

for $k \leq n$. Let $k \in \{1, 2, \dots, n\}$ and $\epsilon > 0$ be sufficiently small compared with k . Then, there exists $\alpha_{k,\epsilon} \in \mathcal{A}$ such that for any $\alpha \geq \alpha_{k,\epsilon}$,

$$(1) \text{ for each } \lambda \in \{\lambda_1, \dots, \lambda_{k-1}\} \text{ with } \lambda < \lambda_k,$$

$$E_\alpha([\lambda - \epsilon, \lambda + \epsilon])L_\alpha^2 = \langle \psi_i^\alpha \mid p_\lambda \leq i \leq q_\lambda \rangle,$$

$$\text{where } p_\lambda = \min\{i \in \mathbb{N} \mid \lambda_i = \lambda\} \text{ and } q_\lambda = \max\{i \in \mathbb{N} \mid \lambda_i = \lambda\} ;$$

$$(2) \text{ for each } i = 1, \dots, k-1 \text{ with } \lambda_i < \lambda_{i+1},$$

$$E_\alpha([\lambda_i + \epsilon, \lambda_{i+1} - \epsilon])L_\alpha^2 = \{0\}.$$

Conclusion Let $a, b \in \mathbb{R}^+ \setminus \sigma(A)$ two given real numbers such that $a < b$, then from what precedes we have for α large enough

$$E_\alpha([a, b])L_\alpha^2 = \langle \psi_k^\alpha \mid k = 1, \dots, n \text{ with } a < \lambda_k \leq b \rangle.$$

Thus $n_\alpha([a, b])$ coincides with the number k such that $a < \lambda_k \leq b$, in other words $n([a, b])$. $\square$

The proof of theorem 20 is the same as above, but defining the Λ_k^α with the help of the φ_k^α .

References

- [BD98] **A. Braides** and **A. Defranceschi**. *Homogenization of Multiple Integrals*. Oxford Science Publications, 1998.
- [Ber86] **P. H. Berard**. *Spectral Geometry: Direct and Inverse Problems*. Number 1207 in Lecture Notes Math. Springer Verlag, 1986.
- [BI94] **D. Burago** and **S. Ivanov**. Riemannian tori without conjugate points are flat. *GAFA*, 4(3):259–269, 1994.
- [BI95] **D. Burago** and **S. Ivanov**. On asymptotic volume of tori. *GAFA*, 5(5):800–808, 1995.
- [BLP78] **A. Bensousan**, **J.-L. Lions**, and **G. Papanicolaou**. *Asymptotic analysis for periodic structures*. Studies in mathematics and its applications. North Holland, 1978.
- [Bow58] **F. Bowman**. *Introduction to Bessel Functions*. Dover, 1958.
- [Bur92] **D. Burago**. Periodic metrics. *Advances in soviet mathematics*, 9:205–210, 1992.
- [Cha94] **I. Chavel**. *Riemannian geometry : a modern introduction*. Cambridge tracts in mathematics. Cambridge University Press, 1994.
- [DZ00] **G. Duro** and **E. Zuazua**. Large time behavior for convection-diffusion equations in $\mathbb{R}^n$ with periodic coefficients. *Journal of Differential Equation*, 167:275–315, 2000.
- [Fed69] **H. Federer**. *Geometric Measure Theory*. Springer Verlag, 1969.
- [Fuk90] **Fukaya**. Hausdorff convergence of Riemann manifold and its application. In **T. Ochiai**, editor, *Recent topic in Differential and Analytic Geometry*, volume 18-I of *Advanced Studies in Pure Mathematics*, pages 143–238. 1990.
- [GHL90] **S. Gallot**, **D. Hulin**, and **J. Lafontaine**. *Riemannian geometry*. Universitext. Springer-Verlag, second edition, 1990.
- [KKO97] **A. Kasue**, **H. Kumura**, and **Y. Ogura**. Convergence of heat kernels on a compact manifold. *Kyushu J. Math*, 51:453–524, 1997.
- [KS] **K. Kuwae** and **T. Shioya**. Convergence of spectral structures: a functional analytic theory and its applications to spectral geometry. prepublication.
- [KS00] **M. Kotani** and **T. Sunada**. Albanese maps and off diagonal long time asymptotics for the heat kernel. *Commun. Math. Phys.*, 209:633–670, 2000.
- [Laf74] **J. Lafontaine**. Sur le volume de la variété de jacobi d’une variété riemannienne. *C. R. Acad. Sc. Paris*, 278:1519–1522, 1974.
- [Mas93] **Dal Maso**. *An Introduction to Γ -convergence*. Birkhäuser, 1993.
- [Mos94] **U. Mosco**. Composite media and asymptotic dirichlet forms. *J. Funct. Anal.*, 123(2):368–421, 1994.

- [MS86] **V. D. Milman** and **G. Schechtman**. *Asymptotic Theory of finite dimensional Normed Spaces*. Number 1200 in Lecture Notes Math. Springer Verlag, 1986.
- [Pan82] **P. Pansu**. *Géométrie du groupe de Heisenberg*. Thèse de docteur 3ème cycle, Université Paris VII, 1982.
- [Pan99] **P. Pansu**. Profil isopérimétrique, métriques périodiques et formes d'équilibre des cristaux. prépublication d'orsay, 1999.
- [Rud91] **W. Rudin**. *Functional Analysis*. International series in Pure and Applied Mathematics. Mc Graw-Hill, second edition, 1991.
- [Ver01] **C. Vernicos**. *Spectres asymptotiques des nilvariétés graduées*. Thèse de doctorat, Université Grenoble I, Joseph Fourier, 2001.
- [ZKON79] **V.V. Zhikov**, **S.M. Kozlov**, **O.A. Oleinik**, and **Kha T'en Ngoan**. Averaging and g-convergence of differential operators. *Russian Math. Surveys*, 34(5):69–147, 1979.

Constantin Vernicos

Université de Neuchâtel

Institut de Mathématiques

11, rue Emile Argand

2007 Neuchâtel

Switzerland

mail: `Constantin.Vernicos@unine.ch`